

(80122)

Medical Benefit		Effective Date: 04/01/13	Next Review Date: 11/17
Preauthorization	Yes	Review Dates: 04/07, 05/08, 01/10, 01/11, 01/12, 01/13, 01/14, 11/14, 11/15, 11/16	

Preauthorization is required and must be obtained through Case Management.

The following Protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> • With a hemoglobinopathy • With a bone marrow failure syndrome • With a primary immunodeficiency • With an inherited metabolic diseases • With a genetic disorder affecting skeletal tissue 	Interventions of interest are: <ul style="list-style-type: none"> • Allogeneic hematopoietic stem cell transplantation 	Comparators of interest are: <ul style="list-style-type: none"> • Standard care 	Relevant outcomes include: <ul style="list-style-type: none"> • Overall survival • Disease-specific survival • Symptoms • Change in disease status • Morbid events • Functional outcomes • Quality of life • Hospitalizations • Medication use • Resource utilization • Treatment-related mortality • Treatment-related morbidity

Description

A number of inherited and acquired conditions have the potential for severe and/or progressive disease. For some conditions, allogeneic hematopoietic stem cell transplantation (allo-HSCT) has been used to alter the natural history of the disease or potentially offer a cure.

Summary of Evidence

The evidence for allo-HSCT in select individuals who have a hemoglobinopathy, bone marrow failure syndrome, primary immunodeficiency, inherited metabolic syndrome diseases, or a genetic disorder affecting skeletal tissue includes mostly case series, case reports, and registry data. Relevant outcomes are overall survival, disease-specific survival, symptoms, change in disease status, morbid events, functional outcomes, quality of life, hospitalizations, medication use, resource utilization, and treatment-related mortality and morbidity. The evidence has shown that, for most of these disorders, there is a demonstrable improvement in overall survival and other disease-specific outcomes. The exception has been with the use of allo-HSCT in the inherited metabolic diseases Hunter, Sanfilippo, and Morquio syndromes. Allo-HSCT is likely to improve health outcomes in select patients with certain inherited and acquired diseases.

Policy

Allogeneic hematopoietic stem cell transplantation is considered **medically necessary** for selected patients with the following disorders.

Hemoglobinopathies

- Sickle cell anemia for children or young adults with either a history of prior stroke or at increased risk of stroke or end-organ damage
- Homozygous beta-thalassemia (i.e., thalassemia major)

Bone marrow failure syndromes

- Aplastic anemia including hereditary (including Fanconi anemia, dyskeratosis congenita, Shwachman-Diamond, Diamond-Blackfan) or acquired (e.g., secondary to drug or toxin exposure) forms.

Primary immunodeficiencies

- Absent or defective T-cell function (e.g., severe combined immunodeficiency, Wiskott-Aldrich syndrome, X-linked lymphoproliferative syndrome)
- Absent or defective natural killer function (e.g., Chediak-Higashi syndrome)
- Absent or defective neutrophil function (e.g., Kostmann syndrome, chronic granulomatous disease, leukocyte adhesion defect)

(See policy guideline #1.)

Inherited metabolic disease

- Lysosomal and peroxisomal storage disorders *except* Hunter, Sanfilippo, and Morquio syndromes

(See policy guideline #2.)

Genetic disorders affecting skeletal tissue

- Infantile malignant osteopetrosis (Albers-Schonberg disease or marble bone disease)

Policy Guidelines

Individual transplant facilities may have their own *additional* requirements or protocols that must be met in order for the patient to be eligible for a transplant at **their** facility.

1. The following lists the immunodeficiencies that have been successfully treated by allogeneic hematopoietic stem-cell transplantation (allo-HSCT) (Gennery & Cant et al, 2008).

Lymphocyte Immunodeficiencies

Adenosine deaminase deficiency

Artemis deficiency

Calcium channel deficiency

CD 40 ligand deficiency

Cernunnos/X-linked lymphoproliferative disease deficiency

CHARGE syndrome with immune deficiency

Common gamma chain deficiency
Deficiencies in CD45, CD3, CD8
DiGeorge syndrome
DNA ligase IV deficiency syndrome
Interleukin-7 receptor alpha deficiency
Janus-associated kinase 3 (JAK3) deficiency
Major histocompatibility class II deficiency
Omenn syndrome
Purine nucleoside phosphorylase deficiency
Recombinase-activating gene (RAG) 1/2 deficiency
Reticular dysgenesis
Winged helix deficiency
Wiskott-Aldrich syndrome
X-linked lymphoproliferative disease
Zeta-chain-associated protein-70 (ZAP-70) deficiency

Phagocytic Deficiencies

Chédiak-Higashi syndrome
Chronic granulomatous disease
Griscelli syndrome type 2
Hemophagocytic lymphohistiocytosis
Interferon-gamma receptor deficiencies
Leukocyte adhesion deficiency
Severe congenital neutropenias
Shwachman-Diamond syndrome

Other Immunodeficiencies

Autoimmune lymphoproliferative syndrome
Cartilage hair hypoplasia
CD25 deficiency
Hyper IgD and IgE syndromes
ICF syndrome
IPEX syndrome
NEMO deficiency
NF- κ B inhibitor, alpha (I κ B-alpha) deficiency

Nijmegen breakage syndrome

2. In the inherited metabolic disorders, allo-HSCT has been proven effective in some cases of Hurler, Maroteaux-Lamy, and Sly syndromes, childhood onset cerebral X-linked adrenoleukodystrophy, globoid-cell leukodystrophy, metachromatic leukodystrophy, alpha-mannosidosis, and aspartylglucosaminuria. Allogeneic HSCT is possibly effective for fucosidosis, Gaucher types 1 and 3, Farber lipogranulomatosis, galactosialidosis, GM₁, gangliosidosis, mucopolysaccharidosis II (I-cell disease), multiple sulfatase deficiency, Niemann-Pick, neuronal ceroid lipofuscinosis, sialidosis, and Wolman disease. Allogeneic HSCT has not been effective in Hunter, Sanfilippo, or Morquio syndromes. (Mehta, 2004).

The experience with reduced-intensity conditioning (RIC) and allogeneic HSCT for the diseases listed in this Protocol has been limited to small numbers of patients, and have yielded mixed results, depending upon the disease category. In general, the results have been most promising in the bone marrow failure syndromes and primary immunodeficiencies. In the hemoglobinopathies, success has been hampered by difficulties with high rates of graft rejection, and in adult patients, severe graft versus host disease (GVHD). Several Phase II/III trials are ongoing examining the role of this type of transplant for these diseases.

Medicare Advantage

If a transplant is needed, we arrange to have the transplant center review and decide whether the patient is an appropriate candidate for the transplant.

Background

Hematopoietic stem cell transplantation (HSCT) refers to a procedure in which hematopoietic stem cells are infused to restore bone marrow function in patients who receive bone-marrow-toxic doses of cytotoxic drugs with or without whole-body radiotherapy. Allo-HSCT refers to the use of hematopoietic progenitor cells obtained from a donor. They can be harvested from bone marrow, peripheral blood, or umbilical cord blood and placenta shortly after delivery of neonates. Cord blood is discussed in greater detail in the Placental and Umbilical Cord Blood as a Source of Stem Cells Protocol.

Immunologic compatibility between infused hematopoietic stem cells and the recipient is a critical factor for achieving a good outcome with allo-HSCT. Compatibility is established by typing of human leukocyte antigen (HLA) using cellular, serologic, or molecular techniques. HLA refers to the tissue type expressed at the HLA-A, -B, and -DR (antigen-D related) loci on each arm of chromosome 6. Depending on the disease being treated, an acceptable donor will match the patient at all or most of the HLA loci (with the exception of umbilical cord blood).

Preparative Conditioning for Allogeneic HSCT

The conventional practice of allo-HSCT involves administration of cytotoxic agents (e.g., cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to destroy endogenous hematopoietic capability in the recipient. Reduced-intensity conditioning (RIC) refers to chemotherapy regimens that seek to reduce adverse effects secondary to bone marrow toxicity. These regimens partially eradicate the patient's hematopoietic ability, thereby allowing for relatively prompt hematopoietic recovery. Patients who undergo RIC with allo-HSCT initially demonstrate donor cell engraftment and bone marrow mixed chimerism. Most will subsequently convert to full-donor chimerism. A number of different cytotoxic regimens, with or without radiotherapy, may be used for RIC allotransplantation. They represent a continuum in their intensity, from almost totally myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition.

Genetic Diseases and Acquired Anemias

Hemoglobinopathies

The thalassemias result from mutations in the globin genes, resulting in reduced or absent hemoglobin production, thereby reducing oxygen delivery. The supportive treatment of β -thalassemia major requires life-long red blood cell transfusions that lead to progressive iron overload and the potential for organ damage and impaired cardiac, hepatic, and endocrine function. The only definitive cure for thalassemia is to correct the genetic defect with allo-HSCT.

Sickle cell disease is caused by a single amino acid substitution in the beta chain of hemoglobin and, unlike thalassemia major, has a variable course of clinical severity.¹ Sickle cell disease typically manifests clinically with anemia, severe painful crises, acute chest syndrome, stroke, chronic pulmonary and renal dysfunction, growth retardation, neurologic deficits, and premature death. The mean age of death for patients with sickle cell disease has been demonstrated as 42 years for men and 48 for women. Three major therapeutic options are available: chronic blood transfusions, hydroxyurea, and allo-HSCT, the latter being the only possibility for cure.¹

Bone Marrow Failure Syndromes

Aplastic anemia in children is rare and is most often idiopathic and less commonly, due to a hereditary disorder. Inherited syndromes include Fanconi anemia, a rare, autosomal recessive disease characterized by genomic instability, with congenital abnormalities, chromosome breakage, cancer susceptibility, and progressive bone marrow failure leading to pancytopenia and severe aplastic anemia. Frequently, this disease terminates in a myelodysplastic syndrome or acute myeloid leukemia (AML). Most patients with Fanconi anemia succumb to the complications of severe aplastic anemia, leukemia, or solid tumors, with a median survival of 30 years of age.² In Fanconi anemia, HSCT is currently the only treatment that definitively restores normal hematopoiesis. Excellent results have been observed with the use of HLA-matched sibling allogeneic HSCT, with cure of the marrow failure and amelioration of the risk of leukemia.²

Dyskeratosis congenita is characterized by marked telomere dysregulation with clinical features of reticulated skin hyperpigmentation, nail dystrophy, and oral leukoplakia.³ Early mortality is associated with bone marrow failure, infections, pulmonary complications, or malignancy.³

Mutations affecting ribosome assembly and function are associated with Shwachman-Diamond syndrome and Diamond-Blackfan anemia.³ Shwachman-Diamond has clinical features that include pancreatic exocrine insufficiency, skeletal abnormalities, and cytopenias, with some patients developing aplastic anemia. As with other bone marrow failure syndromes, patients are at increased risk of myelodysplastic syndrome and malignant transformation, especially AML. Diamond-Blackfan anemia is characterized by absent or decreased erythroid precursors in the bone marrow, with 30% of patients also having a variety of physical anomalies.³

Primary Immunodeficiencies

The primary immunodeficiencies are a genetically heterogeneous group of diseases that affect distinct components of the immune system. More than 120 gene defects have been described, causing more than 150 disease phenotypes.⁴ The most severe defects (collectively known as severe combined immunodeficiency [SCID]) cause an absence or dysfunction of T lymphocytes and sometimes B lymphocytes and natural killer cells.⁴ Without treatment, patients with SCID usually die by 12 to 18 months of age. With supportive care, including prophylactic medication, the lifespan of these patients can be prolonged, but long-term outlook is still poor, with many dying from infectious or inflammatory complications or malignancy by early adulthood.⁴ Bone marrow transplantation is the only definitive cure, and the treatment of choice for SCID and other primary immunodeficiencies, including Wiskott-Aldrich syndrome and congenital defects of neutrophil function.⁵

Inherited Metabolic Diseases

Lysosomal storage disorders consist of many different rare diseases caused by a single gene defect, and most are inherited as an autosomal recessive trait.⁶ Lysosomal storage disorders are caused by specific enzyme deficiencies that result in defective lysosomal acid hydrolysis of endogenous macromolecules that subsequently accumulate as a toxic substance. Peroxisomal storage disorders arise due to a defect in a membrane transporter protein that leads to defects in the metabolism of long-chain fatty acids. Lysosomal storage disorders and peroxisomal storage disorders affect multiple organ systems, including the central and peripheral nervous systems. These disorders are progressive and often fatal in childhood due to both the accumulation of toxic substrate and a deficiency of the product of the enzyme reaction.⁶ Hurler syndrome usually leads to premature death by five years of age.

Exogenous enzyme replacement therapy is available for a limited number of the inherited metabolic diseases; however, these drugs do not cross the blood-brain barrier, which results in ineffective treatment of the central nervous system. Stem cell transplantation provides a constant source of enzyme replacement from the engrafted donor cells, which are not impeded by the blood-brain barrier.⁶ The donor-derived cells can migrate and engraft in many organ systems, giving rise to different types of cells (e.g., microglial cells in the brain and Kupffer cells in the liver).⁶

Allo-HSCT has been primarily used to treat the inherited metabolic diseases that belong to the lysosomal and peroxisomal storage disorders, as listed in Table 1.⁶ The first stem cell transplant for an inherited metabolic disease was performed in 1980 in a patient with Hurler syndrome. Since that time, more than 1000 transplants have been performed worldwide.⁶

Table 1. Lysosomal and Peroxisomal Storage Disorders

Category	Diagnosis	Other Names
Mucopolysaccharidosis (MPS)	MPS I	Hurler, Scheie, H-S
	MPS II	Hunter
	MPS III A-D	Sanfilippo A-D
	MPS IV A-B	Morquio A-B
	MPS VI	Maroteaux-Lamy
	MPS VII	Sly
Sphingolipidosis	Fabry	
	Farber	Lipogranulomatosis
	Gaucher types 1 and 3	
	GM ₁ gangliosidosis	
	Niemann-Pick disease A and B	
	Tay-Sachs disease	
	Sandhoff disease	
	Globoid cell leukodystrophy	Krabbe disease
Metachromatic leukodystrophy	MLD	
Glycoproteinosis	Aspartylglucosaminuria	
	Fucosidosis	
	Alpha-mannosidosis	
	Beta-mannosidosis	
	Mucopolipidosis III and IV	Sialidosis
Other lipidoses	Niemann-Pick disease C	
	Wolman disease	
	Ceroid lipofuscinosis	Type III-Batten disease
Glycogen storage	GSD type II	Pompe
Multiple enzyme deficiency	Galactosialidosis	
	Mucopolipidosis type II	I-cell disease

Category	Diagnosis	Other Names
Lysosomal transport defects	Cystinosis Sialic acid storage disease Salla disease	
Peroxisomal storage disorders	Adrenoleukodystrophy Adrenomyeloneuropathy	ALD AMN

Infantile Malignant Osteopetrosis

Osteopetrosis is a condition caused by defects in osteoclast development and/or function. The osteoclast (the cell that functions in the breakdown and resorption of bone tissue) is known to be part of the hematopoietic family and shares a common progenitor with the macrophage in the bone marrow.⁷ Osteopetrosis is a heterogeneous group of heritable disorders, resulting in several different types of variable severity. The most severely affected patients are those with infantile malignant osteopetrosis (Albers-Schonberg disease or marble bone disease). Patients with infantile malignant osteopetrosis suffer from dense bone, including a heavy head with frontal bossing, exophthalmos, blindness by approximately six months of age, and severe hematologic malfunction with bone marrow failure.⁷ Seventy percent of these patients die before the age of six years, often of recurrent infections.⁷ HSCT is the only curative therapy for this fatal disease.

HSCT for autoimmune disease, such as rheumatoid arthritis or multiple sclerosis, is considered separately in the Hematopoietic Stem-Cell Transplantation for Autoimmune Diseases Protocol.

Regulatory Status

The U.S. Food and Drug Administration regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under the Code of Federal Regulation title 21, parts 1270 and 1271.⁸ Hematopoietic stem cells are included in these regulations.

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this Protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

References

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Bhatia M, Walters MC. Hematopoietic cell transplantation for thalassemia and sickle cell disease: past, present and future. *Bone Marrow Transplant.* Jan 2008; 41(2):109-117. PMID 18059330
2. Mehta P. Hematopoietic stem cell transplantation for inherited bone marrow failure syndromes. In: Mehta P, ed. *Pediatric Stem Cell Transplantation.* Sudbury, MA: Jones and Bartlett Publishers; 2004:281-316.

3. Gluckman E, Wagner JE. Hematopoietic stem cell transplantation in childhood inherited bone marrow failure syndrome. *Bone Marrow Transplant*. Jan 2008; 41(2):127-132. PMID 18084332
4. Gennery AR, Cant AJ. Advances in hematopoietic stem cell transplantation for primary immunodeficiency. *Immunol Allergy Clin North Am*. May 2008; 28(2):439-456, x-xi. PMID 18424341
5. Porta F, Forino C, De Martiis D, et al. Stem cell transplantation for primary immunodeficiencies. *Bone Marrow Transplant*. Jun 2008; 41 Suppl 2(suppl 2):S83-86. PMID 18545252
6. Prasad VK, Kurtzberg J. Emerging trends in transplantation of inherited metabolic diseases. *Bone Marrow Transplant*. Jan 2008; 41(2):99-108. PMID 18176609
7. Askmyr MK, Fasth A, Richter J. Towards a better understanding and new therapeutics of osteopetrosis. *Br J Haematol*. Mar 2008; 140(6):597-609. PMID 18241253
8. Research FaDAFaCfBEa. CBER parts 1270 and 1271. <http://www.fda.gov/BiologicsBloodVaccines/TissueTissueProducts/>. Accessed July 15, 2015.
9. MacMillan ML, Walters MC, Gluckman E. Transplant outcomes in bone marrow failure syndromes and hemoglobinopathies. *Semin Hematol*. Jan 2010; 47(1):37-45. PMID 20109610
10. Smiers FJ, Krishnamurti L, Lucarelli G. Hematopoietic stem cell transplantation for hemoglobinopathies: current practice and emerging trends. *Pediatr Clin North Am*. Feb 2010; 57(1):181-205. PMID 20307718
11. Angelucci E, Matthes-Martin S, Baronciani D, et al. Hematopoietic stem cell transplantation in thalassemia major and sickle cell disease: indications and management recommendations from an international expert panel. *Haematologica*. May 2014; 99(5):811-820. PMID 24790059
12. Mathews V, Srivastava A, Chandy M. Allogeneic stem cell transplantation for thalassemia major. *Hematol Oncol Clin North Am*. Dec 2014; 28(6):1187-1200. PMID 25459187
13. Locatelli F, Kabbara N, Ruggeri A, et al. Outcome of patients with hemoglobinopathies given either cord blood or bone marrow transplantation from an HLA-identical sibling. *Blood*. Aug 8 2013; 122(6):1072-1078. PMID 23692854
14. Mehta P. Hematopoietic stem cell transplantation for hemoglobinopathies. In: Mehta P, ed. *Pediatric Stem Cell Transplantation*. Sudbury, MA: Jones and Bartlett Publishers; 2004:259-279.
15. Mahmoud HK, Elhaddad AM, Fahmy OA, et al. Allogeneic hematopoietic stem cell transplantation for non-malignant hematological disorders. *J Adv Res*. May 2015; 6(3):449-458. PMID 26257943
16. Bernaudin F, Socie G, Kuentz M, et al. Long-term results of related myeloablative stem-cell transplantation to cure sickle cell disease. *Blood*. Oct 1 2007; 110(7):2749-2756. PMID 17606762
17. Walters MC, Patience M, Leisenring W, et al. Bone marrow transplantation for sickle cell disease. *N Engl J Med*. Aug 8 1996; 335(6):369-376. PMID 8663884
18. Walters MC, Storb R, Patience M, et al. Impact of bone marrow transplantation for symptomatic sickle cell disease: an interim report. Multicenter investigation of bone marrow transplantation for sickle cell disease. *Blood*. (6):1918-1924. PMID 10706855
19. Hsieh MM, Fitzhugh CD, Weitzel RP, et al. Nonmyeloablative HLA-matched sibling allogeneic hematopoietic stem cell transplantation for severe sickle cell phenotype. *JAMA*. Jul 2 2014; 312(1):48-56. PMID 25058217
20. Bernardo ME, Piras E, Vacca A, et al. Allogeneic hematopoietic stem cell transplantation in thalassemia major: results of a reduced-toxicity conditioning regimen based on the use of treosulfan. *Blood*. Jul 12 2012; 120(2):473-476. PMID 22645178

21. Anurathapan U, Pakakasama S, Mekjaruskul P, et al. Outcomes of Thalassemia Patients Undergoing Hematopoietic Stem Cell Transplantation by Using a Standard Myeloablative versus a Novel Reduced Toxicity Conditioning Regimen According to a New Risk Stratification. *Biol Blood Marrow Transplant.* Jul 23 2014. PMID 25064743
22. Oringanje C, Nemecek E, Oniyangi O. Hematopoietic stem cell transplantation for people with sickle cell disease. *Cochrane Database Syst Rev.* 2013; 5:CD007001. PMID 23728664
23. Mehta P, Locatelli F, Stary J, et al. Bone marrow transplantation for inherited bone marrow failure syndromes. *Pediatr Clin North Am.* Feb 2010; 57(1):147-170. PMID 20307716
24. Miano M, Dufour C. The diagnosis and treatment of aplastic anemia: a review. *Int J Hematol.* Jun 2015; 101(6):527-535. PMID 25837779
25. Bacigalupo A. Bone marrow transplantation for acquired severe aplastic anemia. *Hematol Oncol Clin North Am.* Dec 2014; 28(6):1145-1155. PMID 25459184
26. Dufour C, Svahn J. Fanconi anaemia: new strategies. *Bone Marrow Transplant.* Jun 2008; 41 Suppl 2 (suppl 2):S90-95. PMID 18545254
27. Zanis-Neto J, Flowers ME, Medeiros CR, et al. Low-dose cyclophosphamide conditioning for haematopoietic cell transplantation from HLA-matched related donors in patients with Fanconi anaemia. *Br J Haematol.* 2005; 130(1):99-106.
28. Wagner JE, Eapen M, MacMillan ML, et al. Unrelated donor bone marrow transplantation for the treatment of Fanconi anemia. *Blood.* Mar 1 2007; 109(5):2256-2262. PMID 17038525
29. Gadalla SM, Sales-Bonfim C, Carreras J, et al. Outcomes of allogeneic hematopoietic cell transplantation in patients with dyskeratosis congenita. *Biol Blood Marrow Transplant.* Aug 2013; 19(8):1238-1243. PMID 23751955
30. Cesaro S, Oneto R, Messina C, et al. Haematopoietic stem cell transplantation for Shwachman-Diamond disease: a study from the European Group for blood and marrow transplantation. *Br J Haematol.* Oct 2005; 131(2):231-236. PMID 16197455
31. Roy V, Perez WS, Eapen M, et al. Bone marrow transplantation for Diamond-Blackfan anemia. *Biol Blood Marrow Transplant.* Aug 2005; 11(8):600-608. PMID 16041310
32. Kim H, Lee JH, Joo YD, et al. A randomized comparison of cyclophosphamide vs. reduced dose cyclophosphamide plus fludarabine for allogeneic hematopoietic cell transplantation in patients with aplastic anemia and hypoplastic myelodysplastic syndrome. *Ann Hematol.* Sep 2012; 91(9):1459-1469. PMID 22526363
33. Dufour C, Pillon M, Socie G, et al. Outcome of aplastic anaemia in children. A study by the severe aplastic anaemia and paediatric disease working parties of the European group blood and bone marrow transplant. *Br J Haematol.* May 2015; 169(4):565-573. PMID 25683884
34. Smith AR, Gross TG, Baker KS. Transplant outcomes for primary immunodeficiency disease. *Semin Hematol.* Jan 2010; 47(1):79-85. PMID 20109615
35. Szabolcs P, Cavazzana-Calvo M, Fischer A, et al. Bone marrow transplantation for primary immunodeficiency diseases. *Pediatr Clin North Am.* Feb 2010; 57(1):207-237. PMID 20307719
36. Ahlin A, Fugelang J, de Boer M, et al. Chronic granulomatous disease-haematopoietic stem cell transplantation versus conventional treatment. *Acta Paediatr.* Nov 2013; 102(11):1087-1094. PMID 23937637

37. Gungor T, Teira P, Slatter M, et al. Reduced-intensity conditioning and HLA-matched haemopoietic stem-cell transplantation in patients with chronic granulomatous disease: a prospective multicentre study. *Lancet*. Feb 1 2014; 383(9915):436-448. PMID 24161820
38. Filipovich A. Hematopoietic cell transplantation for correction of primary immunodeficiencies. *Bone Marrow Transplant*. Aug 2008; 42 Suppl 1(suppl 1):S49-S52. PMID 18724301
39. Hassan A, Booth C, Brightwell A, et al. Outcome of hematopoietic stem cell transplantation for adenosine deaminase-deficient severe combined immunodeficiency. *Blood*. Oct 25 2012; 120(17):3615-3624; quiz 3626. PMID 22791287
40. Filipovich AH, Stone JV, Tomany SC, et al. Impact of donor type on outcome of bone marrow transplantation for Wiskott-Aldrich syndrome: collaborative study of the International Bone Marrow Transplant Registry and the National Marrow Donor Program. *Blood*. Mar 15 2001; 97(6):1598-1603. PMID 11238097
41. Moratto D, Giliani S, Bonfim C, et al. Long-term outcome and lineage-specific chimerism in 194 patients with Wiskott-Aldrich syndrome treated by hematopoietic cell transplantation in the period 1980-2009: an international collaborative study. *Blood*. Aug 11 2011; 118(6):1675-1684. PMID 21659547
42. Marsh RA, Bleesing JJ, Chandrakasan S, et al. Reduced-Intensity Conditioning Hematopoietic Cell Transplantation Is an Effective Treatment for Patients with SLAM-Associated Protein Deficiency/X-linked Lymphoproliferative Disease Type 1. *Biol Blood Marrow Transplant*. Jun 9 2014. PMID 24923536
43. Boelens JJ, Prasad VK, Tolar J, et al. Current international perspectives on hematopoietic stem cell transplantation for inherited metabolic disorders. *Pediatr Clin North Am*. Feb 2010; 57(1):123-145. PMID 20307715
44. Prasad VK, Kurtzberg J. Transplant outcomes in mucopolysaccharidoses. *Semin Hematol*. Jan 2010; 47(1):59-69. PMID 20109613
45. Rovelli AM. The controversial and changing role of haematopoietic cell transplantation for lysosomal storage disorders: an update. *Bone Marrow Transplant*. Jun 2008; 41 Suppl 2(suppl 2):S87-89. PMID 18545253
46. Aldenhoven M, Wynn RF, Orchard PJ, et al. Long-term outcome of Hurler syndrome patients after hematopoietic cell transplantation: an international multicenter study. *Blood*. Mar 26 2015; 125(13):2164-2172. PMID 25624320
47. Mehta P. Metabolic diseases. In: Mehta P, ed. *Pediatric Stem Cell Transplantation*. Sudbury, MA: Jones and Bartlett Publishers; 2004:233-258.
48. Boelens JJ, Wynn RF, O'Meara A, et al. Outcomes of haematopoietic cell transplantation for MPS-1 in Europe: a risk factor analysis for graft failure. *Bone Marrow Transplant*. 2007; 40(3):225-233.
49. Hansen MD, Filipovich AH, Davies SM, et al. Allogeneic hematopoietic cell transplantation (HCT) in Hurler's syndrome using a reduced intensity preparative regimen. *Bone Marrow Transplant*. 2008; 41(4):349-353.
50. Miller WP, Rothman SM, Nascene D, et al. Outcomes after allogeneic hematopoietic cell transplantation for childhood cerebral adrenoleukodystrophy: the largest single-institution cohort report. *Blood*. Aug 18 2011; 118(7):1971-1978. PMID 21586746
51. Mynarek M, Tolar J, Albert MH, et al. Allogeneic hematopoietic SCT for alpha-mannosidosis: an analysis of 17 patients. *Bone Marrow Transplant*. Mar 2012; 47(3):352-359. PMID 21552297
52. Guffon N, Bertrand Y, Forest I, et al. Bone marrow transplantation in children with Hunter syndrome: outcome after 7 to 17 years. *J Pediatr*. May 2009; 154(5):733-737. PMID 19167723

53. Vellodi A, Young E, New M, et al. Bone marrow transplantation for Sanfilippo disease type B. *J Inherit Metab Dis.* 1992; 15(6):911-918. PMID 1293388
54. Bordigoni P, Vidailbet M, Lena M, et al. Bone marrow transplantation for Sanfilippo syndrome. In: Hobbs JR, ed. *Correction of Certain Genetic Diseases by Transplantation.* London: Cogent; 1989:114-119.
55. Steward CG. Hematopoietic stem cell transplantation for osteopetrosis. *Pediatr Clin North Am.* Feb 2010; 57(1):171-180. PMID 20307717
56. Driessen GJ, Gerritsen EJ, Fischer A, et al. Long-term outcome of hematopoietic stem cell transplantation in autosomal recessive osteopetrosis: an EBMT report. *Bone Marrow Transplant.* 2003; 32(7):657-663. PMID
57. Barone A, Lucarelli A, Onofrillo D, et al. Diagnosis and management of acquired aplastic anemia in childhood. Guidelines from the Marrow Failure Study Group of the Pediatric Haemato-Oncology Italian Association (AIEOP). *Blood Cells Mol Dis.* Jun 2015; 55(1):40-47. PMID 25976466