

# Protocol

## Continuous Passive Motion in the Home Setting

(10110)

<b>Medical Benefit</b>		<b>Effective Date:</b> 01/01/12	<b>Next Review Date:</b> 09/17
<b>Preauthorization</b>	Yes	<b>Review Dates:</b> 04/07, 01/08, 11/08, 03/09, 03/10, 09/10, 09/11, 09/12, 09/13, 09/14, 09/15, 09/16	

### ***Preauthorization is required.***

*The following Protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.*

<b>Populations</b>	<b>Interventions</b>	<b>Comparators</b>	<b>Outcomes</b>
Individuals: <ul style="list-style-type: none"><li>• With total knee arthroplasty</li></ul>	Interventions of interest are: <ul style="list-style-type: none"><li>• Continuous passive motion in the home</li></ul>	Comparators of interest are: <ul style="list-style-type: none"><li>• Physical therapy alone</li><li>• Standard of care, if unable to tolerate physical therapy</li></ul>	Relevant outcomes include: <ul style="list-style-type: none"><li>• Symptoms</li><li>• Functional outcomes</li></ul>
Individuals: <ul style="list-style-type: none"><li>• With intra-articular cartilage repair of the knee</li></ul>	Interventions of interest are: <ul style="list-style-type: none"><li>• Continuous passive motion in the home</li></ul>	Comparators of interest are: <ul style="list-style-type: none"><li>• Standard of care</li></ul>	Relevant outcomes include: <ul style="list-style-type: none"><li>• Symptoms</li><li>• Functional outcomes</li></ul>
Individuals: <ul style="list-style-type: none"><li>• With other musculoskeletal conditions requiring physical therapy</li></ul>	Interventions of interest are: <ul style="list-style-type: none"><li>• Continuous passive motion in the home</li></ul>	Comparators of interest are: <ul style="list-style-type: none"><li>• Standard of care</li></ul>	Relevant outcomes include: <ul style="list-style-type: none"><li>• Symptoms</li><li>• Functional outcomes</li></ul>

### **Description**

Continuous passive motion (CPM) devices are used to keep a joint in motion without patient assistance. CPM is being evaluated for treatment and postsurgical rehabilitation of the upper- and lower-limb joints and for a variety of musculoskeletal conditions.

### **Summary of Evidence**

For individuals who have total knee arthroplasty (TKA) who receive continuous passive motion (CPM) in the home, the evidence includes randomized clinical trials (RCTs), case series, and systematic reviews. Relevant outcomes are symptoms and functional outcomes. Early trials generally used CPM in the inpatient setting and are less relevant to today's practice patterns of short hospital stays followed by outpatient rehabilitation. Current postoperative rehabilitation protocols differ considerably from when the largest body of evidence was collected, making it difficult to apply the available evidence to the present situation. For use of CPM after TKA, recent studies have suggested that institutional and home use of CPM has no benefit compared to standard PT. There were no studies evaluating CPM in patients who cannot perform standard PT. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have intra-articular cartilage repair of the knee who receive CPM in the home the evidence includes nonrandomized studies, case series, and studies with nonclinical outcomes (e.g., histology). Relevant outcomes are symptoms and functional outcomes. Systematic reviews of CPM for this indication cite studies reporting better histologic outcomes in patients following CPM. A few studies have reported clinical outcomes, but inadequacies of these studies do not permit conclusions of efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have other musculoskeletal conditions requiring PT who receive CPM in the home, the evidence includes RCTs for some conditions and only case series for others. Relevant outcomes are symptoms and functional outcomes. Three small RCTs of CPM after rotator cuff surgery showed some evidence that CPM after rotator cuff repair of the shoulder improves short-term pain and range of motion; however, the studies were not of high quality, and the small differences in outcomes may not be clinically important. Two of these trials reported short-term improvements in range of motion for patients undergoing CPM, and one reported a short-term reduction in pain. None reported long-term improvements, and there are no reported benefits in functional status. Therefore, the clinical significance of the short-term improvements reported is uncertain. In addition, there is uncertainty about the optimal PT regimen following shoulder surgery such that the optimal comparison for CPM is unclear. For other conditions, RCTs do not exist; case series did not show efficacy of CPM or had important methodologic flaws. The evidence is insufficient to determine the effects of the technology on health outcomes.

### Policy

Use of continuous passive motion (CPM) in the home setting may be considered **medically necessary** as an adjunct to physical therapy in the following situations:

- Under conditions of low postoperative mobility or inability to comply with rehabilitation exercises following a total knee arthroplasty (TKA) or TKA revision. This may include patients with complex regional pain syndrome (reflex sympathetic dystrophy); extensive arthrofibrosis or tendon fibrosis; or physical, mental or behavioral inability to participate in active physical therapy.
- During the non-weight bearing rehabilitation period following intra-articular cartilage repair procedures of the knee (e.g., microfracture, osteochondral grafting, autologous chondrocyte implantation, treatment of osteochondritis dissecans, repair of tibial plateau fractures).

Use of CPM in the home setting for all other indications is considered **not medically necessary**.

### Policy Guidelines

This Protocol only addresses CPM in the home setting (i.e., not the hospital setting).

Following TKA, CPM in the home setting will be allowable for up to 21 days after surgery while patients are immobile or unable to bear weight.

Following intra-articular cartilage repair procedures of the knee, CPM in the home setting will be allowable for up to six weeks during non-weight bearing rehabilitation.

### Medicare Advantage

CPM devices are devices **medically necessary** for patients who have received a total knee replacement.

To qualify for coverage:

- use of the device must commence within two days following surgery
- coverage is limited to that portion of the three-week period following surgery during which the device is used in the patient's home.

### Background

Physical therapy (PT) of joints following surgery focuses both on passive motion to restore mobility and active exercises to restore strength. While passive motion can be administered by a therapist, continuous passive motion (CPM) devices have also been used. CPM is thought to improve recovery by stimulating the healing of articular tissues and the circulation of synovial fluid; reducing local edema; and preventing adhesions, joint stiffness or contractures, or cartilage degeneration. CPM has been most thoroughly investigated in the knee, particularly after TKA or ligamentous or cartilage repair, but acceptance of its use in the knee joint has created interest in extrapolating this experience to other weight-bearing joints (i.e., hip, ankle, metatarsals) and non-weight-bearing joints (i.e., shoulder, elbow, metacarpals, interphalangeal joints). Use of CPM in stroke and burn patients is also being explored.

The device used for the knee moves the joint (e.g., flexion/extension), without patient assistance, continuously for extended periods of time (i.e., up to 24 h/d). An electrical power unit is used to set the variable range of motion (ROM) and speed. The initial settings for ROM are based on a patient's level of comfort and other factors that are assessed intraoperatively. The ROM is increased by 3° to 5° per day, as tolerated. The speed and ROM can be varied, depending on joint stability. The use of the device may be initiated in the immediate postoperative period and then continued at home for a variable period of time.

Over the past 10 to 20 years, hospital lengths of stay have progressively shortened, and in some cases, surgical repair may be done either as an outpatient or with a length of stay of one to two days. As a result, there has been a considerable shift in the rehabilitation regimen, moving from an intensive in-hospital program to a less intensive outpatient program. Therefore, some providers may want patients to continue CPM in the home as a means of duplicating the services offered with a longer (seven-day) hospital stay. The focus of the current review is to examine the literature regarding home use of CPM as it is currently being prescribed postoperatively. The most important comparisons will be treatment outcomes of CPM when used alone or in addition to conventional PT, compared with conventional PT alone.

### Regulatory Status

Continuous passive motion devices are considered class I devices by the U.S. Food and Drug Administration (FDA) and are exempt from 510(k) requirements. This classification does not require submission of clinical data regarding efficacy but only notification of FDA prior to marketing. FDA product code: BXB.

### Related Protocols

Autografts and Allografts in the Treatment of Focal Articular Cartilage Lesions

Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions

---

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are

considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this Protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

## References

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Continuous Passive Motion as an Adjunct to Physical Therapy for Joint Rehabilitation. TEC Assessments. 1997; Volume 12 (Tab 20).
2. McInnes J, Larson MG, Daltroy LH, et al. A controlled evaluation of continuous passive motion in patients undergoing total knee arthroplasty. *JAMA*. Sep 16 1992; 268(11):1423-1428. PMID 1512910
3. Milne S, Brosseau L, Robinson V, et al. Continuous passive motion following total knee arthroplasty. *Cochrane Database Syst Rev*. 2003(2):CD004260. PMID 12804511
4. Brosseau L, Milne S, Wells G, et al. Efficacy of continuous passive motion following total knee arthroplasty: a metaanalysis. *J Rheumatol*. Nov 2004; 31(11):2251-2264. PMID 15517640
5. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev*. 2010(3):CD004260. PMID 20238330
6. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database Syst Rev*. 2014; 2:CD004260. PMID 24500904
7. He ML, Xiao ZM, Lei M, et al. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty. *Cochrane Database Syst Rev*. 2014; 7:CD008207. PMID 25069620
8. Yashar AA, Venn-Watson E, Welsh T, et al. Continuous passive motion with accelerated flexion after total knee arthroplasty. *Clin Orthop Relat Res*. Dec 1997 (345):38-43. PMID 9418619
9. MacDonald SJ, Bourne RB, Rorabeck CH, et al. Prospective randomized clinical trial of continuous passive motion after total knee arthroplasty. *Clin Orthop Relat Res*. Nov 2000 (380):30-35. PMID 11064970
10. Pope RO, Corcoran S, McCaul K, et al. Continuous passive motion after primary total knee arthroplasty. Does it offer any benefits? *J Bone Joint Surg Br*. Nov 1997; 79(6):914-917. PMID 9393903
11. Kumar PJ, McPherson EJ, Dorr LD, et al. Rehabilitation after total knee arthroplasty: a comparison of 2 rehabilitation techniques. *Clin Orthop Relat Res*. Oct 1996 (331):93-101. PMID 8895624
12. Bruun-Olsen V, Heiberg KE, Mengshoel AM. Continuous passive motion as an adjunct to active exercises in early rehabilitation following total knee arthroplasty - a randomized controlled trial. *Disabil Rehabil*. 2009; 31(4):277-283. PMID 18608367
13. Denis M, Moffet H, Caron F, et al. Effectiveness of continuous passive motion and conventional physical therapy after total knee arthroplasty: a randomized clinical trial. *Phys Ther*. Feb 2006; 86(2):174-185. PMID 16445331
14. Leach W, Reid J, Murphy F. Continuous passive motion following total knee replacement: a prospective randomized trial with follow-up to 1 year. *Knee Surg Sports Traumatol Arthrosc*. Oct 2006; 14(10):922-926. PMID 16489477

15. Boese CK, Weis M, Phillips T, et al. The efficacy of continuous passive motion after total knee arthroplasty: a comparison of three protocols. *J Arthroplasty*. Jun 2014; 29(6):1158-1162. PMID 24412145
16. Herbold JA, Bonistall K, Blackburn M, et al. Randomized controlled trial of the effectiveness of continuous passive motion after total knee replacement. *Arch Phys Med Rehabil*. Jul 2014; 95(7):1240-1245. PMID 24685389
17. Herbold JA, Bonistall K, Blackburn M. Effectiveness of continuous passive motion in an inpatient rehabilitation hospital after total knee replacement: a matched cohort study. *PM R*. Oct 2012; 4(10):719-725. PMID 22959052
18. Chen B, Zimmerman JR, Soulen L, et al. Continuous passive motion after total knee arthroplasty: a prospective study. *Am J Phys Med Rehabil*. Sep-Oct 2000; 79(5):421-426. PMID 10994883
19. Worland RL, Arredondo J, Angles F, et al. Home continuous passive motion machine versus professional physical therapy following total knee replacement. *J Arthroplasty*. Oct 1998; 13(7):784-787. PMID 9802665
20. Lenssen TA, van Steyn MJ, Crijs YH, et al. Effectiveness of prolonged use of continuous passive motion (CPM), as an adjunct to physiotherapy, after total knee arthroplasty. *BMC Musculoskelet Disord*. 2008; 9:60. PMID 18442423
21. Browne JE, Anderson AF, Arciero R, et al. Clinical outcome of autologous chondrocyte implantation at 5 years in US subjects. *Clin Orthop Relat Res*. Jul 2005 (436):237-245. PMID 15995447
22. Farr J. Autologous chondrocyte implantation improves patellofemoral cartilage treatment outcomes. *Clin Orthop Relat Res*. Oct 2007; 463:187-194. PMID 17960681
23. Rosenberger RE, Gomoll AH, Bryant T, et al. Repair of large chondral defects of the knee with autologous chondrocyte implantation in patients 45 years or older. *Am J Sports Med*. Dec 2008; 36(12):2336-2344. PMID 18725654
24. Nugent-Derfus GE, Takara T, O'Neill J K, et al. Continuous passive motion applied to whole joints stimulates chondrocyte biosynthesis of PRG4. *Osteoarthritis Cartilage*. May 2007; 15(5):566-574. PMID 17157538
25. Salter RB. The biologic concept of continuous passive motion of synovial joints. The first 18 years of basic research and its clinical application. *Clin Orthop Relat Res*. May 1989 (242):12-25. PMID 2650945
26. Fazalare JA, Griesser MJ, Siston RA, et al. The use of continuous passive motion following knee cartilage defect surgery: a systematic review. *Orthopedics*. Dec 2010; 33(12):878. PMID 21162503
27. Howard JS, Mattacola CG, Romine SE, et al. Continuous passive motion, early weight bearing, and active motion following knee articular cartilage repair: evidence for clinical practice. *Cartilage*. Oct 2010; 1(4):276-286. PMID 26069559
28. Hill AD, Palmer MJ, Tanner SL, et al. Use of continuous passive motion in the postoperative treatment of intra-articular knee fractures. *J Bone Joint Surg Am*. Jul 16 2014; 96(14):e118. PMID 25031380
29. Wright RW, Preston E, Fleming BC, et al. A systematic review of anterior cruciate ligament reconstruction rehabilitation: part I: continuous passive motion, early weight bearing, postoperative bracing, and home-based rehabilitation. *J Knee Surg*. Jul 2008; 21(3):217-224. PMID 18686484
30. Du Plessis M, Eksteen E, Jenneker A, et al. The effectiveness of continuous passive motion on range of motion, pain and muscle strength following rotator cuff repair: a systematic review. *Clin Rehabil*. Apr 2011; 25(4):291-302. PMID 20943710
31. Lastayo PC, Wright T, Jaffe R, et al. Continuous passive motion after repair of the rotator cuff. A prospective outcome study. *J Bone Joint Surg Am*. Jul 1998; 80(7):1002-1011. PMID 9698005

32. Raab MG, Rzeszutko D, O'Connor W, et al. Early results of continuous passive motion after rotator cuff repair: a prospective, randomized, blinded, controlled study. *Am J Orthop (Belle Mead NJ)*. Mar 1996; 25(3):214-220. PMID 8775698
33. Garofalo R, Conti M, Notarnicola A, et al. Effects of one-month continuous passive motion after arthroscopic rotator cuff repair: results at 1-year follow-up of a prospective randomized study. *Musculoskelet Surg*. May 2010; 94 Suppl 1:S79-83. PMID 20383685
34. Simkin PA, de Lateur BJ, Alquist AD, et al. Continuous passive motion for osteoarthritis of the hip: a pilot study. *J Rheumatol*. Sep 1999; 26(9):1987-1991. PMID 10493681
35. Dundar U, Toktas H, Cakir T, et al. Continuous passive motion provides good pain control in patients with adhesive capsulitis. *Int J Rehabil Res*. Sep 2009; 32(3):193-198. PMID 19011582
36. Lynch D, Ferraro M, Krol J, et al. Continuous passive motion improves shoulder joint integrity following stroke. *Clin Rehabil*. Sep 2005; 19(6):594-599. PMID 16180594
37. Lindenhovius AL, van de Luitgaarden K, Ring D, et al. Open elbow contracture release: postoperative management with and without continuous passive motion. *J Hand Surg Am*. May-Jun 2009; 34(5):858-865. PMID 19362791
38. Ring D, Simmons BP, Hayes M. Continuous passive motion following metacarpophalangeal joint arthroplasty. *J Hand Surg Am*. May 1998; 23(3):505-511. PMID 9620192
39. Schwartz DA, Chafetz R. Continuous passive motion after tenolysis in hand therapy patients: a retrospective study. *J Hand Ther*. Jul-Sep 2008; 21(3):261-266; quiz 267. PMID 18652971
40. Zeifang F, Carstens C, Schneider S, et al. Continuous passive motion versus immobilisation in a cast after surgical treatment of idiopathic club foot in infants: a prospective, blinded, randomised, clinical study. *J Bone Joint Surg Br*. Dec 2005; 87(12):1663-1665. PMID 16326882
41. Kasten P, Geiger F, Zeifang F, et al. Compliance with continuous passive movement is low after surgical treatment of idiopathic club foot in infants: a prospective, double-blinded clinical study. *J Bone Joint Surg Br*. Mar 2007; 89(3):375-377. PMID 17356153
42. Gavish L, Barzilay Y, Koren C, et al. Novel continuous passive motion device for self-treatment of chronic lower back pain: a randomised controlled study. *Physiotherapy*. Mar 2015; 101(1):75-81. PMID 25280603
43. American Academy of Orthopaedic Surgeons. Surgical management of osteoarthritis of the knee. Evidence-based clinical practice guideline. 2015; [http://www.aaos.org/uploadedFiles/PreProduction/Quality/Guidelines\\_and\\_Reviews/guidelines/SMOAK%20CPG\\_4.22.2016.pdf](http://www.aaos.org/uploadedFiles/PreProduction/Quality/Guidelines_and_Reviews/guidelines/SMOAK%20CPG_4.22.2016.pdf). Accessed July 8, 2016.
44. Postel JM, Thoumie P, Missaoui B, et al. Continuous passive motion compared with intermittent mobilization after total knee arthroplasty. Elaboration of French clinical practice guidelines. *Ann Readapt Med Phys*. May 2007; 50(4):244-257. PMID 17412445
45. Center for Medicare and Medicaid. National Coverage Decision (NCD) for Durable Medical Equipment Reference List (280.1). 5/5/2005: <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=190&bc=AgAAQAAAAAAAA&ncdver=2>. Accessed February, 2016.