

Protocol

Endovascular Therapies for Extracranial Vertebral Artery Disease

(701148)

Medical Benefit		Effective Date: 07/01/15	Next Review Date: 05/19
Preauthorization	No	Review Dates: 05/15, 05/16, 05/17, 05/18	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: • With extracranial vertebral artery stenosis	Interventions of interest are: • Percutaneous transluminal angioplasty with or without stent implantation	Comparators of interest are: • Medical management	Relevant outcomes include: • Overall survival • Symptoms • Morbid events • Treatment-related mortality • Treatment-related morbidity
Individuals: • With extracranial vertebral artery aneurysm(s)	Interventions of interest are: • Percutaneous transluminal angioplasty with stent implantation	Comparators of interest are: • Observation • Surgical treatment	Relevant outcomes include: • Overall survival • Symptoms • Morbid events • Treatment-related mortality • Treatment-related morbidity
Individuals: • With extracranial vertebral artery dissection(s)	Interventions of interest are: • Percutaneous transluminal angioplasty with stent implantation	Comparators of interest are: • Observation • Surgical treatment	Relevant outcomes include: • Overall survival • Symptoms • Morbid events • Treatment-related mortality • Treatment-related morbidity
Individuals: • With extracranial vertebral artery arteriovenous fistula(e)	Interventions of interest are: • Percutaneous transluminal angioplasty with stent implantation	Comparators of interest are: • Observation • Surgical treatment	Relevant outcomes include: • Overall survival • Symptoms • Morbid events • Treatment-related mortality • Treatment-related morbidity

DESCRIPTION

Vertebral artery diseases, including atherosclerotic stenosis, dissections, and aneurysms, can lead to ischemia of the posterior cerebral circulation. Conventional management of extracranial vertebral artery diseases may include medical therapy, including antiplatelet or anticoagulant medications, medications to reduce athero-

sclerotic disease risk (e.g., statins), and/or surgical revascularization. Endovascular therapies have been investigated as an alternative to conventional management.

SUMMARY OF EVIDENCE

ANGIOPLASTY WITH OR WITHOUT STENTING

For individuals who have extracranial vertebral artery stenosis who receive percutaneous transluminal angioplasty with or without stent implantation, the evidence includes a phase 2 randomized controlled trial (RCT). Relevant outcomes are overall survival, symptoms, morbid events, and treatment-related mortality and morbidity. The phase 2 RCT, the Vertebral Artery Stenting Trial (VAST), found no advantage for endovascular intervention compared to best medical therapy alone, with a periprocedural adverse event rate of 5% for the invasive procedures. A larger phase 3 trial comparing endovascular therapy to medical therapy for vertebral artery stenosis is ongoing, although the lack of benefit of endovascular therapy demonstrated in VAST raises questions about the need for a phase 3 trial. Evidence from noncomparative studies has shown that vertebral artery stenting can be performed with high rates of technical success and low periprocedural morbidity and mortality, and that vessel patency can be achieved in a high percentage of cases. However, long-term follow-up has demonstrated high rates of in-stent stenosis. The evidence is insufficient to determine the effects of the technology on health outcomes.

ANGIOPLASTY WITH STENTING

For individuals who have extracranial vertebral artery aneurysm(s), dissection(s), or arteriovenous (AV) fistula(e) who receive percutaneous transluminal angioplasty with stent implantation, the evidence includes small case series and reports. Relevant outcomes are overall survival, symptoms, morbid events, and treatment-related mortality and morbidity. The available evidence has indicated that endovascular therapy for extracranial vertebral artery disorders other than stenosis is feasible and may be associated with favorable outcomes. However, given the lack of data comparing endovascular therapies to alternatives, the evidence is insufficient to determine whether endovascular therapy for extracranial vertebral artery aneurysms, dissections, or AV fistulae improves the net health outcome. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Endovascular therapy, including percutaneous transluminal angioplasty with or without stenting, is considered **investigational** for the management of extracranial vertebral artery disease.

POLICY GUIDELINES

The extracranial vertebral artery is considered to be segments V1-V3 of the vertebral artery from its origin at the subclavian artery until it crosses the dura mater.

BACKGROUND

VERTEBROBASILAR CIRCULATION ISCHEMIA

Ischemia of the vertebrobasilar or posterior circulation accounts for about 20% of all strokes. Posterior circulation strokes may arise from occlusion of the innominate and subclavian arteries, the extracranial vertebral arteries, or the intracranial vertebral, basilar, or posterior cerebral arteries. Compared with carotid artery

disease, relatively little is known about the true prevalence of specific causes of posterior circulation strokes, particularly the prevalence of vertebral artery disease. Reports from one stroke registry have estimated that, in 9% of cases, posterior circulation strokes are due to stenosis of the proximal vertebral artery.¹ Patients who experience strokes or transient ischemic attacks of the vertebrobasilar circulation face a 25% to 35% risk of stroke within the subsequent five years. In particular, the presence of vertebral artery stenosis increases the 90-day risk of recurrent stroke by about four-fold.

Relevant Clinical Anatomy and Pathophysiology

Large artery disease of the posterior circulation may be due to atherosclerosis (stenosis), embolism, dissection, or aneurysms. In about a third of cases, posterior circulation strokes are due to stenosis of the extracranial vertebral arteries or the intracranial vertebral, basilar, and posterior cerebral arteries. The proximal portion of the vertebral artery in the neck is the most common location of atherosclerotic stenosis in the posterior circulation. Dissection of the extracranial or intracranial vertebral arteries may also cause posterior circulation ischemia. By contrast, posterior cerebral artery ischemic events are more likely to be secondary to embolism from more proximal vessels.

The vertebral artery is divided into four segments, V1 through V4, of which segments V1, V2, and V3 are extracranial. V1 originates at the subclavian artery and extends to the C5 or C6 vertebrae; V2 crosses the bony canal of the transverse foramina from C2 to C5; V3 starts as the artery exits the transverse foramina at C2 and ends as the vessel crosses the dura mater and becomes an intracranial vessel. The most proximal segment (V1) is the most common location for atherosclerotic occlusive disease to occur, while arterial dissections are most likely to involve the extracranial vertebral artery just before the vessel crosses the dura mater. Compared with the carotid circulation, the vertebral artery system is more likely to be associated with anatomic variants, including a unilateral artery.

Atherosclerotic disease of the vertebral artery is associated with conventional risk factors for cerebrovascular disease. However, risk factors and the underlying pathophysiology of vertebral artery dissection and aneurysms differ. Extracranial vertebral artery aneurysms and dissections are most often secondary to trauma, particularly those with excessive rotation, distraction, or flexion/extension, or iatrogenic injury, such as during cervical spine surgeries. Spontaneous vertebral artery dissections are rare, and in many cases are associated with connective tissue disorders, including Ehlers-Danlos syndrome type IV, Marfan syndrome, autosomal dominant polycystic kidney disease, and osteogenesis imperfecta type I.²

Management of Extracranial Vertebral Artery Disease

The optimal management of occlusive extracranial vertebral artery disease is not well-defined. Medical treatment with antiplatelet or anticoagulant medications is a mainstay of therapy to reduce stroke risk. Medical therapy also typically involves risk reduction for classical cardiovascular risk factors. However, no randomized trials have compared specific antiplatelet or anticoagulant regimens.

Surgical revascularization may be used for vertebral artery atherosclerotic disease, but open surgical repair is considered technically challenging due to poor access to the vessel origin. Surgical repair may involve vertebral endarterectomy, bypass grafting, or transposition of the vertebral artery, usually to the common or internal carotid artery. Moderately sized, single-center case series of surgical vertebral artery repair from 2012 and 2013 have reported overall survival rates of 91% and 77% at three and six years postoperatively, and arterial patency rates of 80% after one year of follow-up.^{3,4} Surgical revascularization may be used when symptomatic vertebral artery stenosis is not responsive to medical therapy, particularly when bilateral vertebral artery stenosis is present or when unilateral stenosis is present in the presence of an occluded or hypoplastic contralateral vertebral artery. Surgical revascularization may also be considered in patients with concomitant symptomatic carotid and vertebral disease who do not have relief from vertebrobasilar ischemia after carotid revascularization.

The management of extracranial vertebral artery aneurysms or dissections is controversial due to uncertainty about the risk of thromboembolic events associated with aneurysms and dissections. Antiplatelet therapy is typically used; surgical repair, which may include vertebral bypass, external carotid autograft, and vertebral artery transposition to the internal carotid artery, or endovascular treatment with stent placement or coil embolization, may also be used.

Given the technical difficulties related to surgically accessing the extracranial vertebral artery, endovascular therapies have been investigated for extracranial vertebral artery disease. Endovascular therapy may consist of percutaneous transluminal angioplasty, with or without stent implantation.

REGULATORY STATUS

Currently, no endovascular therapies have been approved by the U.S. Food and Drug Administration (FDA) specifically for treatment of extracranial vertebral artery disease.

Various stents, approved for use in the carotid or coronary circulation, have been used for extracranial vertebral artery disease. These stents may be self- or balloon-expandable.

Two devices have been approved by FDA through the humanitarian device exemption process for *intracranial* atherosclerotic disease. This form of FDA approval is available for devices used to treat conditions with an incidence of 4000 or less per year; FDA only requires data showing “probable safety and effectiveness.” Devices with their labeled indications are as follows:

1. Neurolink System® (Guidant, Santa Clara, CA). “The Neurolink system is indicated for the treatment of patients with recurrent intracranial stroke attributable to atherosclerotic disease refractory to medical therapy in intracranial vessels ranging from 2.5 to 4.5 mm in diameter with $\geq 50\%$ stenosis and that are accessible to the stent system.”
2. Wingspan™ Stent System (Boston Scientific, Fremont, CA). “The Wingspan Stent System with Gateway PTA Balloon Catheter is indicated for use in improving cerebral artery lumen diameter in patients with intracranial atherosclerotic disease, refractory to medical therapy, in intracranial vessels with $\geq 50\%$ stenosis that are accessible to the system.”

RELATED PROTOCOLS

Endovascular Procedures for Intracranial Arterial Disease (Atherosclerosis and Aneurysms)

Extracranial Carotid Artery Stenting

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Gulli G, Marquardt L, Rothwell PM, et al. Stroke risk after posterior circulation stroke/transient ischemic attack and its relationship to site of vertebrobasilar stenosis: pooled data analysis from prospective studies. *Stroke*. Mar 2013; 44(3):598-604. PMID 23386676
2. Morasch MD, Phade SV, Naughton P, et al. Primary extracranial vertebral artery aneurysms. *Ann Vasc Surg*. May 2013; 27(4):418-423. PMID 23540677
3. Coleman DM, Obi A, Criado E, et al. Contemporary outcomes after distal vertebral reconstruction. *J Vasc Surg*. Jul 2013; 58(1):152-157. PMID 23478503
4. Ramirez CA, Febrer G, Gaudric J, et al. Open repair of vertebral artery: a 7-year single-center report. *Ann Vasc Surg*. Jan 2012; 26(1):79-85. PMID 22176877
5. Compter A, van der Worp HB, Schonewille WJ, et al. Stenting versus medical treatment in patients with symptomatic vertebral artery stenosis: a randomised open-label phase 2 trial. *Lancet Neurol*. Jun 2015; 14(6):606-614. PMID 25908089
6. Antoniou GA, Murray D, Georgiadis GS, et al. Percutaneous transluminal angioplasty and stenting in patients with proximal vertebral artery stenosis. *J Vasc Surg*. Apr 2012; 55(4):1167-1177. PMID 22206680
7. Coward LJ, McCabe DJ, Ederle J, et al. Long-term outcome after angioplasty and stenting for symptomatic vertebral artery stenosis compared with medical treatment in the Carotid And Vertebral Artery Transluminal Angioplasty Study (CAVATAS): a randomized trial. *Stroke*. May 2007; 38(5):1526-1530. PMID 17395869
8. Karameshev A, Schroth G, Mordasini P, et al. Long-term outcome of symptomatic severe ostial vertebral artery stenosis (OVAS). *Neuroradiology*. May 2010; 52(5):371-379. PMID 20148328
9. Stayman AN, Nogueira RG, Gupta R. A systematic review of stenting and angioplasty of symptomatic extracranial vertebral artery stenosis. *Stroke*. Aug 2011; 42(8):2212-2216. PMID 21700936
10. Coward LJ, Featherstone RL, Brown MM. Percutaneous transluminal angioplasty and stenting for vertebral artery stenosis. *Cochrane Database Syst Rev*. 2005(2):CD000516. PMID 15846607
11. Kikuchi T, Ishii A, Nakahara I, et al. Japanese Registry of Neuroendovascular Therapy: extracranial stenotic diseases except for internal carotid artery stenosis. *Neurol Med Chir (Tokyo)*. 2014; 54(1):40-45. PMID 24257542
12. Sun X, Ma N, Wang B, et al. The long term results of vertebral artery ostium stenting in a single center. *J Neurointerv Surg*. Dec 2015; 7(12):888-891. PMID 25332411
13. Mohammadian R, Sharifipour E, Mansourizadeh R, et al. Angioplasty and stenting of symptomatic vertebral artery stenosis. Clinical and angiographic follow-up of 206 cases from Northwest Iran. *Neuroradiol J*. Aug 2013; 26(4):454-463. PMID 24007733
14. Hatano T, Tsukahara T, Miyakoshi A, et al. Stent placement for atherosclerotic stenosis of the vertebral artery ostium: angiographic and clinical outcomes in 117 consecutive patients. *Neurosurgery*. Jan 2011; 68(1):108-116; discussion 116. PMID 21099720
15. Pham MH, Rahme RJ, Arnaout O, et al. Endovascular stenting of extracranial carotid and vertebral artery dissections: a systematic review of the literature. *Neurosurgery*. Apr 2011; 68(4):856-866; discussion 866. PMID 21242839
16. Badve MS, Henderson RD, O'Sullivan JD, et al. Vertebrobasilar dissections: case series comparing patients with and without dissecting aneurysms. *J Clin Neurosci*. Nov 2014; 21(11):2028-2030. PMID 24913932
17. Horowitz MB, Miller G, 3rd, Meyer Y, et al. Use of intravascular stents in the treatment of internal carotid and extracranial vertebral artery pseudoaneurysms. *AJNR Am J Neuroradiol*. Apr 1996; 17(4):693-696. PMID 8730189
18. Felber S, Henkes H, Weber W, et al. Treatment of extracranial and intracranial aneurysms and arteriovenous fistulae using stent grafts. *Neurosurgery*. Sep 2004; 55(3):631-638; discussion 638-639. PMID 15335430

19. Herrera DA, Vargas SA, Dublin AB. Endovascular treatment of traumatic injuries of the vertebral artery. *AJNR Am J Neuroradiol*. Sep 2008; 29(8):1585-1589. PMID 18499790
20. Ambekar S, Sharma M, Smith D, et al. Successful treatment of iatrogenic vertebral pseudoaneurysm using pipeline embolization device. *Case Rep Vasc Med*. 2014; 2014:341748. PMID 25276469
21. Jang HJ, Oh SY, Shim YS, et al. Endovascular treatment of symptomatic high-flow vertebral arteriovenous fistula as a complication after c1 screw insertion. *J Korean Neurosurg Soc*. Oct 2014; 56(4):348-352. PMID 25371787
22. Shang EK, Fairman RM, Foley PJ, et al. Endovascular treatment of a symptomatic extracranial vertebral artery aneurysm. *J Vasc Surg*. Nov 2013; 58(5):1391-1393. PMID 23561429
23. Takahashi S, Katayama K, Tatsugawa T, et al. A successful hybrid repair for vertebral arteriovenous fistula with extracranial vertebral artery aneurysm. *Ann Vasc Surg*. Jan 2015; 29(1):126 e125-128. PMID 25304908
24. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Jul 2014; 45(7):2160-2236. PMID 24788967
25. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *Circulation*. Jul 26 2011; 124(4):e54-130. PMID 21282504
26. European Stroke Organisation, Tendera M, Aboyans V, et al. ESC Guidelines on the diagnosis and treatment of peripheral artery diseases: Document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries: the Task Force on the Diagnosis and Treatment of Peripheral Artery Diseases of the European Society of Cardiology (ESC). *Eur Heart J*. Nov 2011; 32(22):2851-2906. PMID 21873417
27. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for Percutaneous Transluminal Angioplasty (PTA) (20.7). 2013; <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=201&ncdver=10&bc=AgAAgAAAAAAAAA%3d%3d&>. Accessed May 7, 2016.
28. National Government Services, Inc. (Primary Geographic Jurisdiction - Illinois, New York - Entire State, Connecticut, Massachusetts, Maine, New Hampshire, Rhode Island, Vermont, Wisconsin, Minnesota) Local Coverage Determination (LCD): Category III CPT® Codes (L33392), Revision Effective Date for services performed on or after 01/01/2018.