

(20196)

Medical Benefit		Effective Date: 04/01/16	Next Review Date: 01/21
Preauthorization	No	Review Dates: 01/15, 01/16, 01/17, 01/18, 01/19, 01/20	

Preauthorization is not required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> With signs and/or symptoms of autonomic nervous system dysfunction 	Interventions of interest are: <ul style="list-style-type: none"> Autonomic nervous system testing 	Comparators of interest are: <ul style="list-style-type: none"> Clinical workup without autonomic nervous system testing 	Relevant outcomes include: <ul style="list-style-type: none"> Test accuracy Symptoms Functional outcomes Quality of life

DESCRIPTION

The autonomic nervous system (ANS) controls physiologic processes that are not under conscious control. ANS testing consists of a battery of tests intended to evaluate the integrity and function of the ANS. These tests are intended as adjuncts to the clinical examination in the diagnosis of ANS disorders.

SUMMARY OF EVIDENCE

For individuals who have signs and symptoms of ANS dysfunction who receive ANS testing, the evidence includes studies of diagnostic accuracy. Relevant outcomes are test accuracy, symptoms, functional outcomes, and quality of life. The evidence base is limited. There is a lack of a criterion standard for determining autonomic dysfunction, which limits the ability to perform high-quality research on diagnostic accuracy. Also, numerous tests are used in various conditions, making it difficult to determine values for the overall diagnostic accuracy of a battery of tests. Scattered reports of diagnostic accuracy are available for certain tests, most commonly in the diabetic population, but these reports do not specify estimates of accuracy for the entire battery of tests. Reported sensitivities and specificities are high for patients with clinically defined distal symmetric polyneuropathy using a symptom-based score as a reference standard, but these estimates are likely biased by study designs that used patients with clinically diagnosed disease and a control group of healthy volunteers. The evidence is insufficient to determine the effects of the technology on health outcomes.

Clinical input obtained in 2014 has strongly supported the use of ANS testing as a diagnostic aid in situations of suspected ANS disorders. ANS testing should be performed in the setting of a dedicated ANS testing laboratory. Portable, automated testing intended for office use has not been sufficiently validated and has a greater potential to lead to erroneous results. Also, despite the limitations in the evidence base, ANS tests provide information that cannot be obtained by other methods, given the limitations of clinical examination. Information

from ANS testing can improve diagnostic accuracy in some patients, and in others may obviate the need for further diagnostic testing.

POLICY

Autonomic nervous system testing, consisting of a battery of tests in several domains (see Policy Guidelines) may be considered **medically necessary** when the following criteria are met:

- Signs and/or symptoms of autonomic dysfunction are present; AND
- A definitive diagnosis cannot be made from clinical examination and routine laboratory testing alone; AND
- Diagnosis of the suspected autonomic disorder will lead to a change in management or will eliminate the need for further testing.

Autonomic nervous system testing is considered **investigational** in all other situations when criteria are not met, including but not limited to the evaluation of the following conditions:

- chronic fatigue syndrome
- fibromyalgia
- anxiety and other psychologic disorders
- sleep apnea
- allergic conditions
- hypertension
- screening of asymptomatic individuals
- monitoring progression of disease or response to treatment.

Autonomic nervous system testing using portable automated devices is considered **investigational** for all indications (see Policy Guidelines).

POLICY GUIDELINES

Although there is no standard battery of tests for ANS testing, a full battery of testing generally consists of individual tests in three categories.

- Cardiovagal function (heart rate variability, heart rate response to deep breathing and Valsalva maneuver)
- Vasomotor adrenergic function (blood pressure response to standing, Valsalva, maneuver, hand grip, tilt table testing)
- Sudomotor function (Quantitative Sudomotor Axon Reflex Test, quantitative sensory test, Thermoregulatory Sweat Test, silastic sweat imprint, sympathetic skin response, electrochemical sweat conductance).

At least one test in each category is usually performed. More than one test from a category will often be included in a battery of tests, but the incremental value of using multiple tests in a category is unknown.

There is little evidence on the comparative accuracy of different ANS tests, but the following tests are generally considered to have uncertain value in ANS testing:

- Pupillography

- Pupil edge light cycle
- Gastric emptying tests
- Cold pressor test
- Quantitative direct and indirect testing of sudomotor function test
- Plasma catecholamine levels
- Skin vasomotor testing
- The ANSAR® test.

ANS testing should be performed in a dedicated ANS testing laboratory. Testing in a dedicated laboratory should be performed under closely controlled conditions, and results should be interpreted by an individual with expertise in ANS testing. Testing using automated devices with results interpreted by computer software has not been validated and thus has the potential to lead to erroneous results.

MEDICARE ADVANTAGE

For Medicare Advantage autonomic function testing (ANS) is considered **medically necessary** when used as a diagnostic tool to evaluate symptoms indicative of vasomotor instability and the ANS testing is directed at establishing a more accurate or definitive diagnosis or contributing to clinically useful and relevant medical decision making for one of the following indications:

- To diagnose the presence of autonomic neuropathy in a patient with signs or symptoms suggesting a progressive autonomic neuropathy.
- To evaluate the severity and distribution of a diagnosed progressive autonomic neuropathy.
- To differentiate the diagnosis between certain complicated variants of syncope from other causes of loss of consciousness.
- To evaluate inadequate response to beta blockade in vasodepressor syncope.
- To evaluate distressing symptoms in a patient with a clinical picture suspicious for distal small fiber neuropathy in order to diagnose the condition.
- To differentiate the cause of postural tachycardia syndrome.
- To evaluate change in type, distribution or severity of autonomic deficits in patients with autonomic failure.
- To evaluate the response to treatment in patients with autonomic failure who demonstrate a change in clinical exam.
- To diagnose axonal neuropathy or suspected autonomic neuropathy in the symptomatic patient.
- To evaluate and treat patients with recurrent unexplained syncope or demonstrate autonomic failure, after more common causes have been excluded by other standard testing.

The following indications are considered **not medically necessary** and will not be covered:

- Screening patients without signs or symptoms of autonomic dysfunction, including patients with diabetes, hepatic or renal disease.
- Testing for the sole purpose of monitoring disease intensity or treatment efficacy in diabetes, hepatic or renal disease.

- Testing results that are not used in clinical decision-making or patient management.
- Testing performed by physicians who do not have evidence of training, and expertise to perform and interpret these tests.
- General professional standards with FDA clearance apply for all equipment used in ANS testing.
- Testing with ANSAR ANX 3.0 or a similar machine is considered investigational for screening.

BACKGROUND

AUTONOMIC NERVOUS SYSTEM

The autonomic nervous system (ANS) has a primary role in controlling physiologic processes not generally under conscious control. They include heart rate, respirations, gastrointestinal (GI) motility, thermal regulation, bladder control, and sexual function.^{1,2} The ANS is a complex neural regulatory network that consists of two complementary systems that work to maintain homeostasis: the sympathetic and the parasympathetic systems. The sympathetic nervous system is responsible for arousal, and sympathetic stimulation leads to increased pulse, increased blood pressure (BP), increased sweating, decreased GI motility, and an increase in other glandular exocrine secretions. This is typically understood as the “fight or flight” response. Activation of the parasympathetic nervous system will mostly have the opposite effects; BP and pulse will decrease, GI motility increases, and there will be a decrease in sweating and other glandular secretions.

ANS DISORDERS

ANS disorders, also called dysautonomias, are heterogeneous in etiology, clinical symptoms, and severity. ANS disorders can be limited and focal, such as patients with isolated neurocardiogenic syncope or idiopathic palmar hyperhidrosis. At the other extreme, some ANS disorders can be widespread and severely disabling, such as multiple systems atrophy, which leads to widespread and severe autonomic failure.

Symptoms of autonomic disorders can vary, based on the etiology and location of dysfunction. Cardiovascular manifestations are often prominent. Involvement of the cardiovascular system causes abnormalities in heart rate control and vascular dynamics.³ Orthostatic hypotension and other manifestations of BP lability can occur, causing weakness, dizziness, and syncope. Resting tachycardia and an inability to appropriately increase heart rate in response to exertion leads to exercise intolerance. There is a two to three fold higher incidence of major cardiac events in patients with diabetic autonomic neuropathy (myocardial infarction, heart failure, resuscitation from ventricular arrhythmia, angina, or the need for revascularization).⁴ There is also an increase in sudden cardiac death and overall mortality for these patients.³

Many other organ systems can be affected by autonomic neuropathy. Involvement of the bladder can lead to incomplete emptying, resulting in urinary retention and possible overflow incontinence. GI involvement is commonly manifested as gastroparesis, which is defined as slowed gastric emptying and can cause nausea, vomiting, and a decreased tolerance for solid food and large meals. Constipation may also occur if the lower GI tract is involved. Impairment of sexual function in males can manifest as erectile dysfunction and ejaculatory failure. Dysfunction of thermal regulation and sweating can lead to anhidrosis and heat intolerance. Paradoxically, excessive sweating can also occur as a compensatory mechanism in unaffected regions.⁵

A classification of the different types of autonomic dysfunction, adapted from Freeman (2005)⁵ and Macdougall and McLeod (1996),⁶ can be made as follows:

- Diabetic autonomic neuropathy
- Amyloid neuropathy
- Immune-mediated neuropathy

- Rheumatoid arthritis
- Systemic lupus erythematosus
- Sjögren syndrome
- Paraneoplastic neuropathy
- Inflammatory neuropathy
 - Guillain-Barré syndrome
 - Chronic inflammatory demyelinating polyneuropathy
 - Crohn disease
 - Ulcerative colitis
- Hereditary autonomic neuropathies
- Autonomic neuropathy secondary to infectious disease
 - HIV disease
 - Lyme disease
 - Chagas disease
 - Diphtheria
 - Leprosy
- Acute and subacute idiopathic autonomic neuropathy
- Toxic neuropathies.

Other chronic diseases may involve an ANS imbalance, without outright dysfunction of the nerves themselves. Approximately 40% of individuals with essential hypertension will show evidence of excess sympathetic activity.⁷ Sympathetic overactivity is also a prominent feature of generalized anxiety, panic disorder, and some types of depression, as well as certain cardiac disorders such as chronic heart failure. These types of ANS imbalances are not usually classified as ANS disorders.

Treatment

Much of the treatment for autonomic disorders is nonpharmacologic and supportive. However, there are specific actions that can improve symptoms in patients with specific deficits. For patients with orthostatic hypotension, this involves adequate intake of fluids and salt, moving to an upright position slowly and deliberately, use of lower-extremity compression stockings, and keeping the head of the bed elevated four to six inches (10-15 cm).¹ In severe cases, treatment with medications that promote salt retention, such as fludrocortisone, is often prescribed. Patients with symptoms of hyperhidrosis may benefit from cooling devices and potent antiperspirants such as Drysol, and patients with decreased tearing and dry mucous membranes can use over-the-counter artificial tears or other artificial moisturizers.¹

ANS Testing

ANS testing consists of a battery of tests. Any single test may be performed individually, or the entire battery of tests may be ordered. Individual components of testing may include cardiovascular function testing, sudomotor function, salivation testing, and tilt table testing.

Cardiovagal Function Testing

Beat-to-beat variability in the heart rate can be measured at rest, or in response to provocative measures, such as deep breathing or the Valsalva maneuver. Reduced, or absent, heart rate variability is a sign of autonomic dysfunction.⁸

Baroreflex sensitivity is measured by examining the change in pulse and heart rate variability in response to changes in BP. A medication such as phenylephrine is given to induce a raise in BP, and baroreflex sensitivity is calculated as the slope of the relation between heart rate variability and BP.⁸

Sudomotor Function (Sweat Testing)

Sweat testing evaluates the structure and function of nerves that regulate the sweat glands.

The Quantitative Sudomotor Axon Reflex Test is an example of a commercially available semiquantitative test of sudomotor function.⁸ The test is performed by placing the color-sensitive paper on the skin, which changes color on contact with sweat. Measurement of the amount of color change is a semiquantitative measure of sudomotor function.

For the silastic sweat imprint, silastic material is placed on the skin, and the sweat droplets form indentations on the silastic surface, allowing quantitation of the degree of sweating present. The Neuropad test is an example of a commercially available silastic sweat imprint.

A more complex approach in some centers is the use of a thermoregulatory laboratory.⁹ This is a closed chamber in which an individual sits for a defined period under tightly controlled temperature and humidity. An indicator dye is brushed on the skin, and it changes color when in contact with sweat. Digital pictures are taken and projected onto anatomic diagrams. Computer processing derives values for a total area of anhidrosis and the percent of anhidrotic areas.

Sympathetic skin response tests use an electric current to stimulate sympathetic nerves. The tests measure the change in electrical resistance, which is altered in the presence of sweat. In general, these tests are considered to be sensitive but have high variability and potential for false-positive results.⁹

A variant of sympathetic skin response testing is electrochemical sweat conductance measured by iontophoresis (e.g., Sudoscan). In this test, a low-level current is used to attract chloride ions from sweat glands. The chloride ions interact with stainless-steel plate electrodes to measure electrochemical resistance.

Salivation Testing

The protocol for salivation testing involves the subject chewing on a pre-weighed gauze for five minutes. At the end of five minutes, the gauze is removed and reweighed to determine the total weight of saliva present.

Tilt Table Testing

Tilt table testing is intended to evaluate for orthostatic intolerance. The patient lies on the table and is strapped in with a foot rest. The table is then inclined to the upright position, with monitoring of the pulse and BP. Symptoms of lightheadedness or syncope in conjunction with changes in pulse or BP constitute a positive test. A provocative medication, such as isoproterenol, can be given to increase the sensitivity of the test.

Composite Autonomic Severity Score

The Composite Autonomic Severity Score, which ranges from zero to 10, is intended to estimate the severity of autonomic dysfunction. Scores are based on self-reported symptoms measured by a standardized symptom survey. Scores of three or less are considered mild, scores of three to seven are considered moderate and scores greater than seven are considered severe.

REGULATORY STATUS

Since 1976, numerous ANS testing devices have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process. Table 1 lists examples.

Table 1. Autonomic Nervous System Test Devices

Device	Manufacturer	Measurement	510(k) No.	Clearance Date	Product Code
ANX 3.0	Ansar Group	Respiration and heart rate variability	K941252	2004	DRT
Sudoscan®	Impeto Medical	Electrochemical sweat conductance	K100233	2010	GZO
ZYTO Hand Cradle	ZYTO Technologies	Galvanic skin response	K111308	2011	GZO
Bodytronic® 200	Bauerfeind	Photoelectric plethysmograph	K123921	2013	JMO
Neuropad®	TRIGOCare	Sudomotor function			

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Klein CM. Evaluation and management of autonomic nervous system disorders. *Semin Neurol.* Apr 2008; 28(2):195-204. PMID 18351521.
2. Gibbons CH, Cheshire WP, Fife TD. American Academy of Neurology Model Coverage Policy: Autonomic Nervous System Testing. 2014; https://www.aan.com/uploadedFiles/Website_Library_Assets/Documents/3.Practice_Management/1.Reimbursement/1.Billing_and_Coding/5.Coverage_Policies/14%20Autonomic%20Testing%20Policy%20v001.pdf. Accessed October 28, 2014.
3. Vinik AI, Ziegler D. Diabetic cardiovascular autonomic neuropathy. *Circulation.* Jan 23 2007;115(3):387-397. PMID 17242296.
4. Valensi P, Sachs RN, Harfouche B, et al. Predictive value of cardiac autonomic neuropathy in diabetic patients with or without silent myocardial ischemia. *Diabetes Care.* Feb 2001;24(2):339-343. PMID 11213889.
5. Freeman R. Autonomic peripheral neuropathy. *Lancet.* Apr 2-8 2005;365(9466):1259-1270. PMID 15811460.
6. McDougall AJ, McLeod JG. Autonomic neuropathy, II: Specific peripheral neuropathies. *J Neurol Sci.* Jun 1996;138(1-2):1-13. PMID 8791232.
7. Goldstein DS, Robertson D, Esler M, et al. Dysautonomias: clinical disorders of the autonomic nervous system. *Ann Intern Med.* Nov 5 2002;137(9):753-763. PMID 12416949.

8. England JD, Gronseth GS, Franklin G, et al. Practice Parameter: evaluation of distal symmetric polyneuropathy: role of autonomic testing, nerve biopsy, and skin biopsy (an evidence-based review). Report of the American Academy of Neurology, American Association of Neuromuscular and Electrodiagnostic Medicine, and American Academy of Physical Medicine and Rehabilitation. *Neurology*. Jan 13 2009;72(2):177-184. PMID 19056667.
9. Low PA. Testing the autonomic nervous system. *Semin Neurol*. Dec 2003;23(4):407-421. PMID 15088262.
10. Bellavere F, Ragazzi E, Chillelli NC, et al. Autonomic testing: which value for each cardiovascular test? An observational study. *Acta Diabetol*. 2019 Jan;56(1):39-43. PMID: 30159748.
11. da Silva AKF, Penachini da Costa de Rezende Barbosa M, Marques Vanderlei F, et al. Application of heart rate variability in diagnosis and prognosis of individuals with diabetes mellitus: systematic review. *Ann Noninvasive Electrocardiol*. May 2016;21(3):223-235. PMID 27226209.
12. Kamenov ZA, Petrova JJ, Christov VG. Diagnosis of diabetic neuropathy using simple somatic and a new autonomic (neuropad) tests in the clinical practice. *Exp Clin Endocrinol Diabetes*. Apr 2010;118(4):226-233. PMID 20200815.
13. Quattrini C, Jeziorska M, Tavakoli M, et al. The Neuropad test: a visual indicator test for human diabetic neuropathy. *Diabetologia*. Jun 2008;51(6):1046-1050. PMID 18368386.
14. Ponirakis G, Petropoulos IN, Fadavi H, et al. The diagnostic accuracy of Neuropad for assessing large and small fibre diabetic neuropathy. *Diabet Med*. Dec 2014;31(12):1673-1680. PMID 24975286.
15. Casellini CM, Parson HK, Richardson MS, et al. Sudoscan, a noninvasive tool for detecting diabetic small fiber neuropathy and autonomic dysfunction. *Diabetes Technol Ther*. Nov 2013;15(11):948-953. PMID 23889506.
16. American Association of Neuromuscular Electrodiagnostic Medicine. Proper performance of autonomic function testing. *Muscle Nerve*. Jan 2017;55(1):3-4. PMID 27786371.
17. American Diabetes Association. Standards of medical care in diabetes--2010. *Diabetes Care*. Jan 2010;33(Suppl 1):S11-61. PMID 20042772.
18. Lahrmann H, Cortelli P, Hilz M, et al. Orthostatic hypotension. In: Gilhus NE, Barnes MP, Brainin M, eds. *European Handbook of Neurological Management: Volume 1, 2nd Edition*. Hoboken, NJ: Wiley-Blackwell; 2011.
19. National Government Services, Inc. (Primary Geographic Jurisdiction 06 & K - Illinois, Minnesota, Wisconsin, Connecticut, New York - Entire State, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont) Local Coverage Determination (LCD): Autonomic Function Testing (L36236), Revision Effective Date for services performed on or after 09/12/2019.