

(70302)

Medical Benefit		Effective Date: 07/01/14	Next Review Date: 05/21
Preauthorization	Yes	Review Dates: 09/09, 09/10, 09/11, 07/12, 05/13, 05/14, 05/15, 05/16, 05/17, 05/18, 05/19, 05/20	

Preauthorization is required and must be obtained through Case Management.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: • With insulin-dependent diabetes	Interventions of interest are: • Pancreas transplant after a kidney transplant	Comparators of interest are: • Insulin therapy	Relevant outcomes include: • Overall survival • Change in disease status • Treatment-related mortality • Treatment-related morbidity
Individuals: • With insulin-dependent diabetes with uremia	Interventions of interest are: • Simultaneous pancreas and kidney transplant	Comparators of interest are: • Insulin therapy	Relevant outcomes include: • Overall survival • Change in disease status • Treatment-related mortality • Treatment-related morbidity
Individuals: • With insulin-dependent diabetes and severe complications	Interventions of interest are: • Pancreas transplant alone	Comparators of interest are: • Insulin therapy	Relevant outcomes include: • Overall survival • Change in disease status • Treatment-related mortality • Treatment-related morbidity
Individuals: • With a prior pancreas transplant who still meet criteria for a pancreas transplant	Interventions of interest are: • Pancreas retransplant	Comparators of interest are: • Insulin therapy	Relevant outcomes include: • Overall survival • Change in disease status • Treatment-related mortality • Treatment-related morbidity

DESCRIPTION

Transplantation of a healthy pancreas is a treatment for patients with insulin-dependent diabetes. Pancreas transplantation can restore glucose control and prevent, halt, or reverse the secondary complications from diabetes.

SUMMARY OF EVIDENCE

For individuals who have insulin-dependent diabetes who receive a pancreas transplant after a kidney trans-

plant, the evidence includes case series and registry studies. The relevant outcomes are overall survival (OS), change in disease status, and treatment-related mortality and morbidity. Data from national and international registries have found relatively high patient survival rates with a pancreas transplant after a kidney transplant (e.g., a three-year survival rate of 93%). A 2012 analysis of data from a single-center found similar patient survival and death-censored pancreas graft survival rates with a pancreas transplant after a kidney transplant or a simultaneous pancreas and kidney (SPK) transplant. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have insulin-dependent diabetes with uremia who receive SPK transplants, the evidence includes registry studies. The relevant outcomes are OS, change in disease status, and treatment-related mortality and morbidity. Data from national and international registries have found relatively high patient survival rates after SPK transplant. A retrospective analysis found a higher survival rate in patients with type 1 diabetes who had an SPK transplant vs. those on a waiting list. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have insulin-dependent diabetes and severe complications who receive pancreas transplant alone, the evidence includes registry studies. The relevant outcomes are OS, change in disease status, and treatment-related mortality and morbidity. Data from international and national registries have found that graft and patient survival rates after pancreas transplant alone have improved over time (e.g., three-year survival of 95%). The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have had a prior pancreas transplant who still meet criteria for a pancreas transplant who receive pancreas retransplantation, the evidence includes case series and registry studies. The relevant outcomes are OS, change in disease status, and treatment-related mortality and morbidity. National data and specific transplant center data have generally found similar graft and patient survival rates after pancreas retransplantation compared with initial transplantation. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

POLICY

Pancreas transplant after a prior kidney transplant may be considered **medically necessary** in patients with insulin-dependent diabetes.

A combined pancreas and kidney transplant may be considered **medically necessary** in insulin-dependent diabetic patients with uremia.

Pancreas transplant alone may be considered **medically necessary** in patients with severely disabling and potentially life-threatening complications due to hypoglycemia unawareness and labile insulin-dependent diabetes that persists in spite of optimal medical management.

Pancreas retransplant after a failed primary pancreas transplant may be considered **medically necessary** in patients who meet criteria for pancreas transplantation.

Pancreas transplant is considered **investigational** in all other situations.

POLICY GUIDELINES

GENERAL

Individual transplant facilities may have their own *additional* requirements or protocols that must be met in order for the patient to be eligible for a transplant at their facility.

Potential contraindications subject to the judgment of the transplant center:

1. Known current malignancy, including metastatic cancer
2. Recent malignancy with high risk of recurrence
3. Untreated systemic infection making immunosuppression unsafe, including chronic infection
4. Other irreversible end-stage disease not attributed to kidney disease
5. History of cancer with a moderate risk of recurrence
6. Systemic disease that could be exacerbated by immunosuppression
7. Psychosocial conditions or chemical dependency affecting ability to adhere to therapy.

PANCREAS SPECIFIC CRITERIA

Candidates for pancreas transplant alone should additionally meet one of the following severity of illness criteria:

- Documentation of severe hypoglycemia unawareness as evidenced by chart notes or emergency department visits; OR
- Documentation of potentially life-threatening labile diabetes as evidenced by chart notes or hospitalization for diabetic ketoacidosis.

Additionally, most pancreas transplant patients will have type 1 diabetes mellitus. Those transplant candidates with type 2 diabetes mellitus, in addition to being insulin-dependent, should also not be obese (body mass index [BMI] should be 32 kg/m² or less). According to International Registry data, in 2010, 7% of pancreas transplant recipients had type 2 diabetes (Gruessner, 2011).

MULTIPLE TRANSPLANT CRITERIA

Although there are no standard guidelines for multiple pancreas transplants, the following information may aid in case review:

- If there is early graft loss resulting from technical factors (e.g., venous thrombosis), a retransplant may generally be performed without substantial additional risk.
- Long-term graft losses may result from chronic rejection, which is associated with increased risk of infection following long-term immunosuppression, and sensitization, which increases the difficulty of finding a negative cross-match. Some transplant centers may wait to allow reconstitution of the immune system before initiating retransplant with an augmented immunosuppression protocol.

MEDICARE ADVANTAGE

If a transplant is needed, we arrange to have the Medicare–approved transplant center review and decide whether the patient is an appropriate candidate for the transplant.

BACKGROUND

Pancreas transplantation occurs in several different scenarios such as (1) a diabetic patient with renal failure who may receive a simultaneous cadaveric pancreas plus kidney transplants; (2) a diabetic patient who may receive a cadaveric or living-related pancreas transplant after a kidney transplantation (pancreas after kidney); or (3) a nonnumeric diabetic patient with specific severely disabling and potentially life-threatening diabetic problems who may receive a pancreas transplant alone. The total number of adult pancreas transplants (pan-

creas and pancreas plus kidney) in the U.S. peaked at 1,484 in 2004 and has since steadily declined.¹ In 2017, 213 received a pancreas transplant alone and 789 simultaneous pancreas plus kidneys were performed in the U.S.¹

According to the International Pancreas Transplant Registry data, the proportion of pancreas transplant recipients worldwide who have type 2 diabetes has increased over time, from 2% in 1995 to 7% in 2010.² In 2010, approximately 8% of simultaneous pancreas plus kidney transplants, 5% of pancreas transplant after kidney transplant, and 1% of a pancreas transplant alone were performed in patients with type 2 diabetes.

REGULATORY STATUS

Small bowel/liver and multivisceral transplantation are surgical procedures and, as such, are not subject to regulation by the U.S. Food and Drug Administration.

The U.S. Food and Drug Administration regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation Title 21, parts 1270 and 1271. Pancreas transplants are included in these regulations.

RELATED PROTOCOLS

Artificial Pancreas Device Systems

Islet Transplantation

Kidney Transplant

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Organ Procurement and Transplantation Network (OPTN). National Data. n.d.; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/>. Accessed August 29, 2019.
2. Gruessner AC. 2011 update on pancreas transplantation: comprehensive trend analysis of 25,000 cases followed up over the course of twenty-four years at the International Pancreas Transplant Registry (IPTR). *Rev Diabet Stud*. Spring 2011;8(1):6-16. PMID 21720668.
3. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Pancreas Transplantation. TEC Assessments. 1998;Volume 13, Tab 7.
4. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Pancreas Retransplantation. TEC Assessments. 2001;Volume 16, Tab 23.

5. Gruessner AC, Gruessner RW. Pancreas Transplantation of US and Non-US Cases from 2005 to 2014 as Reported to the United Network for Organ Sharing (UNOS) and the International Pancreas Transplant Registry (IPTR). *Rev Diabet Stud*. Spring 2016;13(1):35-58. PMID 26982345.
6. Bazerbachi F, Selzner M, Marquez MA, et al. Pancreas-after-kidney versus synchronous pancreas-kidney transplantation: comparison of intermediate-term results. *Transplantation*. Feb 15 2013;95(3):489-494. PMID 23183776.
7. Fridell JA, Mangus RS, Hollinger EF, et al. The case for pancreas after kidney transplantation. *Clin Transplant*. Aug-Sep 2009;23(4):447-453. PMID 19453642.
8. Barlow AD, Saeb-Parsy K, Watson CJE. An analysis of the survival outcomes of simultaneous pancreas and kidney transplantation compared to live donor kidney transplantation in patients with type 1 diabetes: a UK Transplant Registry study. *Transpl Int*. Sep 2017;30(9):884-892. PMID 28319322.
9. van Dellen D, Worthington J, Mitu-Pretorian OM, et al. Mortality in diabetes: pancreas transplantation is associated with significant survival benefit. *Nephrol Dial Transplant*. May 2013;28(5):1315-1322. PMID 23512107.
10. Sampaio MS, Kuo HT, Bunnapradist S. Outcomes of simultaneous pancreas-kidney transplantation in type 2 diabetic recipients. *Clin J Am Soc Nephrol*. May 2011;6(5):1198-1206. PMID 21441123.
11. Scalea JR, Butler CC, Munivenkatappa RB, et al. Pancreas transplant alone as an independent risk factor for the development of renal failure: a retrospective study. *Transplantation*. Dec 27 2008;86(12):1789-1794. PMID 19104423.
12. Gasteiger S, Cardini B, Göbel G, et al. Outcomes of pancreas retransplantation in patients with pancreas graft failure. *Br J Surg*. 2018 Dec;105(13):1816-1824. PMID: 30007018.
13. Buron F, Thaunat O, Demuylder-Mischler S, et al. Pancreas retransplantation: a second chance for diabetic patients? *Transplantation*. Jan 27 2013;95(2):347-352. PMID 23222920.
14. Fridell JA, Mangus RS, Chen JM, et al. Late pancreas retransplantation. *Clin Transplant*. Jan 2015;29(1):1-8. PMID 25284041.
15. Seal J, Selzner M, Laurence J, et al. Outcomes of pancreas retransplantation after simultaneous kidney-pancreas transplantation are comparable to pancreas after kidney transplantation alone. *Transplantation*. Mar 2015;99(3):623-628. PMID 25148379.
16. Rudolph EN, Finger EB, Chandolias N, et al. Outcomes of pancreas retransplantation. *Transplantation*. Feb 2015;99(2):367-374. PMID 25594555.
17. Organ Procurement and Transplantation Network (OPTN). Organ Procurement and Transplantation Network Policies. 2018; https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf. August 29, 2019.
18. Working Party of the British Transplantation Society. Kidney and Pancreas Transplantation in Patients with HIV. Second Edition (Revised). British Transplantation Society Guidelines. Macclesfield, UK: British Transplantation Society; 2017.
19. Siskind E, Maloney C, Akerman M, et al. An analysis of pancreas transplantation outcomes based on age groupings—an update of the UNOS database. *Clin Transplant*. Sep 2014;28(9):990-994. PMID 24954160.
20. Afaneh C, Rich BS, Aull MJ, et al. Pancreas transplantation: does age increase morbidity? *J Transplant*. 2011; 2011:596801. PMID 21766007.
21. Schenker P, Vonend O, Kruger B, et al. Long-term results of pancreas transplantation in patients older than 50 years. *Transpl Int*. Feb 2011;24(2):136-142. PMID 21039944.
22. Gruessner AC, Sutherland DE. Access to pancreas transplantation should not be restricted because of age: invited commentary on Schenker et al [commentary]. *Transpl Int*. Feb 2011;24(2):134-135. PMID 21208293.
23. Organ Procurement and Transplantation Network. Organ Procurement and Transplantation Network (OPTN) Policies. Effective May 23, 2019. https://optn.transplant.hrsa.gov/media/1200/optn_policies.pdf. Accessed July 1, 2019.

24. Centers for Medicare & Medicaid Services (CMS). Medicare-approved transplant programs. 2018; <https://www.cms.gov/Medicare/Provider-Enrollment-and-Certification/CertificationandCompliance/downloads/ApprovedTransplantPrograms.pdf>. Accessed August 29, 2019.
25. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for pancreas Transplants (260.3). 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncdid=107&ver=3>. Accessed August 29, 2019.