

Protocol

Biofeedback as a Treatment of Chronic Pain

(20130)

Medical Benefit		Effective Date: 07/15/07	Next Review Date: 05/21
Preauthorization	No	Review Dates: 02/07, 01/08, 01/09, 01/10, 09/10, 07/11, 07/12, 05/13, 05/14, 05/15, 05/16, 05/17, 05/18, 05/19, 05/20	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none">• With chronic pain	Interventions of interest are: <ul style="list-style-type: none">• Electromyography biofeedback	Comparators of interest are: <ul style="list-style-type: none">• Pharmacologic treatment• Nonpharmacologic treatment	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Functional outcomes• Quality of life• Medication use

DESCRIPTION

Biofeedback is a technique intended to teach patients self-regulation of certain physiologic processes not normally considered to be under voluntary control. Electromyography biofeedback has been evaluated as a method to reduce chronic or recurrent pain of musculoskeletal or psychosomatic origin.

SUMMARY OF EVIDENCE

For individuals who have chronic pain (including low back, knee, neck and shoulder, orofacial, and abdominal pain as well as fibromyalgia, osteoarthritis, systemic lupus erythematosus, and vulvar vestibulitis) who receive biofeedback, the evidence includes multiple randomized controlled trials (RCTs) for different pain syndromes. The relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. The results of these RCTs, some of which were sham-controlled, did not consistently report a benefit for biofeedback. Some RCTs reported improved outcomes with biofeedback, but these improvements were often of uncertain clinical significance or were not durable. Many other RCTs have found that biofeedback did not provide a significantly greater benefit in outcomes when it was used instead of or in addition to other conservative interventions such as exercise. Overall, the available RCTs were limited by small sample sizes and high dropout rates. This evidence base does not permit conclusions about the specific effects of biofeedback beyond the nonspecific effects of sham interventions, nor does it permit conclusions about the contribution of biofeedback beyond that of other conservative treatments for pain. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Biofeedback as a treatment of chronic pain, including but not limited to low back pain, is **investigational**.

BACKGROUND

Treatment for chronic pain is often multimodal and typically includes psychological therapy. Psychological techniques vary but may include cognitive therapy, which teaches subjects the ability to cope with stressful stimuli by attempting to alter negative thought patterns and dysfunctional attitudes, and behavioral approaches to reduce muscle tension and break the pain cycle. Relaxation, using any of a variety of techniques including meditation or mental imagery, is considered a behavioral therapy that may be used alone or as a component of a cognitive-behavioral therapy program. Electromyography biofeedback has also been used for the treatment of chronic pain, on the assumption that the ability to reduce muscle tension will be improved through the feedback of data to the patient regarding the degree of muscle tension. While some consider electromyography biofeedback to be a method used to obtain relaxation, others consider biofeedback to be distinct from other relaxation techniques.

Biofeedback provides physiologic information not normally available to the patient, with a concerted effort employed by the patient to use this feedback to help alter the physiologic process in some specific way. Biofeedback training is done either in individual or group sessions, alone or in combination with other behavioral therapies designed to teach relaxation. A typical program consists of 10 to 20 training sessions of 30 minutes each. Training sessions are performed in a quiet, non-stimulating environment. Patients are instructed to use mental imagery techniques to affect the physiologic variable being monitored, and feedback is provided for the successful alteration of that physiologic parameter in the form of lights or tone, verbal praise, or other auditory or visual stimuli.

REGULATORY STATUS

Since 1976, a large number of biofeedback devices have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process. Food and Drug Administration product code: HCC.

RELATED PROTOCOLS

Biofeedback as a Treatment of Fecal Incontinence or Constipation

Biofeedback as a Treatment of Headache

Biofeedback as a Treatment of Urinary Incontinence in Adults

Biofeedback for Miscellaneous Indications

Neurofeedback

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced

procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Biofeedback. TEC Assessments. 1995;Volume 10:Tab 25.
2. Eccleston C, Williams AC, Morley S. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev*. Apr 15 2009(2):CD007407. PMID 19370688.
3. Eccleston C, Palermo TM, Williams AC, et al. Psychological therapies for the management of chronic and recurrent pain in children and adolescents. *Cochrane Database Syst Rev*. Apr 15 2009(2):CD003968. PMID 19370592.
4. Humphreys PA, Gevirtz RN. Treatment of recurrent abdominal pain: components analysis of four treatment protocols. *J Pediatr Gastroenterol Nutr*. Jul 2000;31(1):47-51. PMID 10896070.
5. Palermo TM, Eccleston C, Lewandowski AS, et al. Randomized controlled trials of psychological therapies for management of chronic pain in children and adolescents: an updated meta-analytic review. *Pain*. Mar 2010; 148(3):387-397. PMID 19910118.
6. Henschke N, Ostelo RW, van Tulder MW, et al. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev*. Jul 07 2010(7):CD002014. PMID 20614428.
7. Kapitza KP, Passie T, Bernateck M, et al. First non-contingent respiratory biofeedback placebo versus contingent biofeedback in patients with chronic low back pain: a randomized, controlled, double-blind trial. *Appl Psychophysiol Biofeedback*. Sep 2010;35(3):207-217. PMID 20237953.
8. Tan G, Rintala DH, Jensen MP, et al. A randomized controlled trial of hypnosis compared with biofeedback for adults with chronic low back pain. *Eur J Pain*. Feb 2015;19(2):271-280. PMID 24934738.
9. Glombiewski JA, Hartwich-Tersek J, Rief W. Two psychological interventions are effective in severely disabled, chronic back pain patients: a randomised controlled trial. *Int J Behav Med*. Jun 2010;17(2):97-107. PMID 19967572.
10. Collins NJ, Bisset LM, Crossley KM, et al. Efficacy of nonsurgical interventions for anterior knee pain: systematic review and meta-analysis of randomized trials. *Sports Med*. Jan 1 2012;42(1):31-49. PMID 22149696.
11. Ma C, Szeto GP, Yan T, et al. Comparing biofeedback with active exercise and passive treatment for the management of work-related neck and shoulder pain: a randomized controlled trial. *Arch Phys Med Rehabil*. Jun 2011;92(6):849-858. PMID 21621660.
12. Ribeiro D, Silva AG. A single session of visual feedback improves range of motion in patients with chronic idiopathic neck pain: A randomized and controlled study. *Musculoskeletal Care*, 2018 Nov 1;17(1). PMID 30378756.
13. Aggarwal VR, Lovell K, Peters S, et al. Psychosocial interventions for the management of chronic orofacial pain. *Cochrane Database Syst Rev*. Nov 09 2011(11):CD008456. PMID 22071849.
14. McNeely ML, Armijo Olivo S, Magee DJ. A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther*. May 2006;86(5):710-725. PMID 16649894.
15. Medicott MS, Harris SR. A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder. *Phys Ther*. Jul 2006;86(7):955-973. PMID 16813476.
16. Weydert JA, Ball TM, Davis MF. Systematic review of treatments for recurrent abdominal pain. *Pediatrics*. Jan 2003;111(1):e1-11. PMID 12509588.

17. Glombiewski JA, Bernardy K, Hauser W. Efficacy of EMG- and EEG-biofeedback in fibromyalgia syndrome: a meta-analysis and a systematic review of randomized controlled trials. *Evid Based Complement Alternat Med*. Oct 2013;2013:962741. PMID 24082911.
18. Buckelew SP, Conway R, Parker J, et al. Biofeedback/relaxation training and exercise interventions for fibromyalgia: a prospective trial. *Arthritis Care Res*. Jun 1998;11(3):196-209. PMID 9782811.
19. van Santen M, Bolwijn P, Verstappen F, et al. A randomized clinical trial comparing fitness and biofeedback training versus basic treatment in patients with fibromyalgia. *J Rheumatol*. Mar 2002;29(3):575-581. PMID 11908576.
20. Babu AS, Mathew E, Danda D, et al. Management of patients with fibromyalgia using biofeedback: a randomized control trial. *Indian J Med Sci*. Aug 2007;61(8):455-461. PMID 17679735.
21. Macfarlane GJ, Paudyal P, Doherty M, et al. A systematic review of evidence for the effectiveness of practitioner-based complementary and alternative therapies in the management of rheumatic diseases: osteoarthritis. *Rheumatology (Oxford)*. Dec 2012;51(12):2224-2233. PMID 22923762.
22. Yilmaz OO, Senocak O, Sahin E, et al. Efficacy of EMG-biofeedback in knee osteoarthritis. *Rheumatol Int*. Aug 20 2010;30:887-892. PMID 19693508.
23. Durmus D, Alayli G, Canturk F. Effects of quadriceps electrical stimulation program on clinical parameters in the patients with knee osteoarthritis. *Clin Rheumatol*. May 2007;26(5):674-678. PMID 16897119.
24. Greco CM, Rudy TE, Manzi S. Effects of a stress-reduction program on psychological function, pain, and physical function of systemic lupus erythematosus patients: a randomized controlled trial. *Arthritis Rheum*. Aug 15 2004;51(4):625-634. PMID 15334437.
25. Bergeron S, Binik YM, Khalife S, et al. A randomized comparison of group cognitive-behavioral therapy, surface electromyographic biofeedback, and vestibulectomy in the treatment of dyspareunia resulting from vulvar vestibulitis. *Pain*. Apr 2001;91(3):297-306. PMID 11275387.
26. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. Apr 04 2017; 166(7):514-530. PMID 28192789.
27. American College of Occupational and Environmental Medicine (ACOEM). Low back disorders. 2016; <https://www.guideline.gov/summaries/summary/50687/low-back-disorders?q=Low+back+disorders>. Accessed September 9, 2019.
28. American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. Apr 2010;112(4):810-833. PMID 20124882.
29. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Biofeedback Therapy (30.1). n.d.; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=41&ncdver=1&bc=AAAAQAAAAA&>. Accessed September 9, 2019.