

Protocol

Dynamic Posturography

(20102)

Medical Benefit		Effective Date: 04/01/13	Next Review Date: 01/20
Preauthorization	No	Review Dates: 05/07, 07/08, 05/09, 03/10, 01/11, 01/12, 01/13, 01/14, 01/15, 01/16, 01/17, 01/18, 01/19	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none">• With suspected balance disorders	Interventions of interest are: <ul style="list-style-type: none">• Dynamic posturography	Comparators of interest are: <ul style="list-style-type: none">• Alternative approach to balance assessment or no balance assessment	Relevant outcomes include: <ul style="list-style-type: none">• Test accuracy• Test validity• Symptoms• Morbid events

DESCRIPTION

Dynamic posturography tests a patient's balance control in situations intended to isolate factors that affect balance in everyday experiences. Posturography provides quantitative information on the degree of imbalance present but is not intended to diagnosis specific types of balance disorders.

SUMMARY OF EVIDENCE

For individuals with suspected balance disorders who receive dynamic posturography, the evidence includes cross-sectional comparisons of results in patients with balance disorders and healthy controls and retrospective case series reporting outcomes for patients assessed with dynamic posturography as part of clinical care. Relevant outcomes are test accuracy and validity, symptoms, and morbid events. There are no generally accepted reference standards for dynamic posturography, which makes it difficult to determine how testing results can be applied to clinical care. There are no studies demonstrating the clinical utility of the test that would lead to changes in management that improve outcomes (e.g., symptoms, function). The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Dynamic posturography is considered **investigational**.

BACKGROUND

BALANCE DISORDERS

Complaints of imbalance are common in older adults and contribute to the risk of falling in this population. Falls are an important cause of death and disability in this population in the United States. Maintenance of balance is a complex physiologic process, requiring the interaction of the vestibular, visual, and proprioceptive/somatosensory system, and central reflex mechanisms. Balance is also influenced by the general health of the patient (i.e., muscle tone, strength, range of motion). Therefore, identifying and treating the underlying balance disorder can be difficult. Commonly used balance function tests (e.g., electronystagmography, rotational chair tests) attempt to measure the extent and site of a vestibular lesion but do not assess the functional ability to maintain balance.

Role in Diagnosis

Dynamic posturography aims to provide quantitative information on a patient's functional ability to maintain balance. The patient, wearing a harness to prevent falls, stands on an enclosed platform surrounded by a visual field. By altering the angle of the platform or shifting the visual field, the test assesses movement coordination and the sensory organization of visual, somatosensory, and vestibular information relevant to postural control. The patient undergoes six different testing situations designed to evaluate the vestibular, visual, and proprioceptive/somatosensory components of balance. In general terms, the test measures an individual's balance (as measured by a force platform to calculate the movement of the patient's center of mass) while visual and somatosensory cues are altered. These tests vary by whether eyes are open or closed, the platform is fixed or sway-referenced, and whether the visual surround is fixed or sway-referenced. Sway-referencing involves making instantaneous computer-aided alterations to the platform or visual surround to coincide with changes in body position produced by sway. The purpose of sway-referencing is to cancel out accurate feedback from somatosensory or visual systems that are normally involved in maintaining balance. In the first three components of the test, the support surface is stable, and visual cues are either present, absent, or sway-referenced. In tests four to six, the support surface is sway-referenced to the individual, and visual cues are either present, absent, or sway-referenced. In tests five and six, the only accurate sensory cues available for balance are vestibular cues. Results of computerized dynamic posturography have been used to determine what type of information (i.e., visual, vestibular, proprioceptive) can and cannot be used to maintain balance. Dynamic posturography cannot be used to localize the site of a lesion.

Posturography tests a patient's balance control in situations intended to isolate factors that affect balance in everyday experiences. Balance can be rapidly assessed qualitatively by asking the patient to maintain a steady stance on a flat or compressible surface (i.e., foam pads) with the eyes open or closed. By closing the eyes, the visual input into balance is eliminated. Use of foam pads eliminates the sensory and proprioceptive cues. Therefore, the only vestibular input is available when standing on a foam pad with eyes closed.

REGULATORY STATUS

In 1985, the NeuroCom EquiTest® (NeuroCom International, Portland, OR; now Clackamas, OR), a dynamic posturography device, was cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process. Other dynamic posturography device makers include Vestibular Technologies (Cheyenne, WY) and Medicateurs (Balma, France). Companies that previously manufactured dynamic posturography devices include Metitur (Jyvaskyla, Finland) and Micromedical Technology (Chatham, IL). Food and Drug Administration product code: LXV.

RELATED PROTOCOL

Vestibular Function Testing

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Dynamic posturography in the assessment of vestibular dysfunction. TEC Assessment. 1996;Volume 11:Tab 11.
2. Fritz NE, Newsome SD, Eloyan A, et al. Longitudinal relationships among posturography and gait measures in multiple sclerosis. *Neurology*. May 19 2015;84(20):2048-2056. PMID 25878185
3. Ferrazzoli D, Fasano A, Maestri R, et al. Balance dysfunction in Parkinson's disease: the role of posturography in developing a rehabilitation program. *Parkinsons Dis*. Oct 2015;2015:520128. PMID 26504611
4. Buatois S, Gueguen R, Gauchard GC, et al. Posturography and risk of recurrent falls in healthy noninstitutionalized persons aged over 65. *Gerontology*. Aug 2006;52(6):345-352. PMID 16905886
5. Girardi M, Konrad HR, Amin M, et al. Predicting fall risks in an elderly population: computer dynamic posturography versus electronystagmography test results. *Laryngoscope*. Sep 2001;111(9):1528-1532. PMID 11568601
6. Sinaki M, Lynn SG. Reducing the risk of falls through proprioceptive dynamic posture training in osteoporotic women with kyphotic posturing: a randomized pilot study. *Am J Phys Med Rehabil*. Apr 2002;81(4):241-246. PMID 11953540
7. Whitney SL, Marchetti GF, Schade AI. The relationship between falls history and computerized dynamic posturography in persons with balance and vestibular disorders. *Arch Phys Med Rehabil*. Mar 2006;87(3):402-407. PMID 16500176
8. Ganesan M, Pasha SA, Pal PK, et al. Direction specific preserved limits of stability in early progressive supranuclear palsy: a dynamic posturographic study. *Gait Posture*. Apr 2012;35(4):625-629. PMID 22225854
9. Lee JM, Koh SB, Chae SW, et al. Postural instability and cognitive dysfunction in early Parkinson's disease. *Can J Neurol Sci*. Jul 2012;39(4):473-482. PMID 22728854
10. Pierchala K, Lachowska M, Morawski K, et al. Sensory Organization Test outcomes in young, older and elderly healthy individuals - preliminary results. *Otolaryngol Pol*. Jul 2012;66(4):274-279. PMID 22890532
11. Biggan JR, Melton F, Horvat MA, et al. Increased load computerized dynamic posturography in prefrail and nonfrail community-dwelling older adults. *J Aging Phys Act*. Jan 2014;22(1):96-102. PMID 23416307
12. Lim KB, Lee HJ. Computerized posturographic measurement in elderly women with unilateral knee osteoarthritis. *Ann Rehabil Med*. Oct 2012;36(5):618-626. PMID 23185725
13. Alahmari KA, Marchetti GF, Sparto PJ, et al. Estimating postural control with the balance rehabilitation unit: measurement consistency, accuracy, validity, and comparison with dynamic posturography. *Arch Phys Med Rehabil*. Jan 2014;95(1):65-73. PMID 24076084
14. Teggi R, Caldirola D, Fabiano B, et al. Rehabilitation after acute vestibular disorders. *J Laryngol Otol*. Apr 2009;123(4):397-402. PMID 18549515

15. Badke MB, Miedaner JA, Shea TA, et al. Effects of vestibular and balance rehabilitation on sensory organization and dizziness handicap. *Ann Otol Rhinol Laryngol*. Jan 2005;114(1 Pt 1):48-54. PMID 15697162
16. Badke MB, Shea TA, Miedaner JA, et al. Outcomes after rehabilitation for adults with balance dysfunction. *Arch Phys Med Rehabil*. Feb 2004;85(2):227-233. PMID 14966706
17. Brown KE, Whitney SL, Marchetti GF, et al. Physical therapy for central vestibular dysfunction. *Arch Phys Med Rehabil*. Jan 2006;87(1):76-81. PMID 16401442
18. Hirsch MA, Toole T, Maitland CG, et al. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch Phys Med Rehabil*. Aug 2003;84(8):1109-1117. PMID 12917847
19. Nocera J, Horvat M, Ray CT. Effects of home-based exercise on postural control and sensory organization in individuals with Parkinson disease. *Parkinsonism Relat Disord*. Dec 2009;15(10):742-745. PMID 19640769
20. Lundin F, Ledin T, Wikkelso C, et al. Postural function in idiopathic normal pressure hydrocephalus before and after shunt surgery: A controlled study using computerized dynamic posturography (EquiTest). *Clin Neurol Neurosurg*. Sep 2013;115(9):1626-1631. PMID 23489444
21. American Academy of Otolaryngology-Head and Neck Surgery. Position Statement: Posturography. 2007 (revised 2014); <http://www.entnet.org/Practice/policyPosturography.cfm>. Accessed January 16, 2018.
22. Bhattacharyya N, Baugh RF, Orvidas L, et al. Clinical practice guideline: benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*. Nov 2008;139(5 Suppl 4):S47-81. PMID 18973840