

(20228)

Medical Benefit		Effective Date: 06/01/19	Next Review Date: 03/21
Preauthorization	Yes	Review Dates: 03/12, 03/13, 03/14, 03/15, 03/16, 03/17, 03/18, 03/19, 03/20	

Preauthorization is required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> Who are asymptomatic with risk for hypertrophic cardiomyopathy because of a positive family history 	Interventions of interest are: <ul style="list-style-type: none"> Testing for a specific hypertrophic cardiomyopathy-related variant identified in affected family member(s) 	Comparators of interest are: <ul style="list-style-type: none"> Clinical management without genetic testing 	Relevant outcomes include: <ul style="list-style-type: none"> Overall survival Test accuracy Test validity Changes in reproductive decision making Symptoms Morbid events
Individuals: <ul style="list-style-type: none"> Who are asymptomatic with risk for hypertrophic cardiomyopathy because of a positive family history 	Interventions of interest are: <ul style="list-style-type: none"> Nonspecific testing for a hypertrophic cardiomyopathy-related variant 	Comparators of interest are: <ul style="list-style-type: none"> Clinical management without genetic testing 	Relevant outcomes include: <ul style="list-style-type: none"> Overall survival Test accuracy Test validity Changes in reproductive decision making Symptoms Morbid events

DESCRIPTION

Familial hypertrophic cardiomyopathy (HCM) is an inherited condition caused by a disease-associated variant in one or more of the cardiac sarcomere genes. HCM is associated with numerous cardiac abnormalities, the most serious of which is sudden cardiac death. Genetic testing for HCM-associated variants is available through a number of commercial laboratories.

SUMMARY OF EVIDENCE

For individuals who are asymptomatic with risk for HCM because of a positive family history who receive testing for a specific HCM-related variant identified in affected family member(s), the evidence includes studies reporting on the clinical validity of testing. Relevant outcomes are overall survival, test accuracy and validity, changes in reproductive decision making, symptoms, and morbid events. For individuals at-risk for HCM (first-degree relatives), genetic testing is most useful when there is a known disease-associated variant in the family. In this

situation, genetic testing will establish the presence or absence of the same variant in a close relative with a high degree of certainty. Presence of the variant indicates that the relative should undergo a cardiac evaluation upon receiving the variant-positive results. If an HCM diagnosis is not made at that time, the patient should be monitored for development of symptoms. Absence of this variant will establish that the individual has not inherited the familial predisposition to HCM and thus has a similar risk of developing HCM as the general population. Such patients will no longer need ongoing surveillance for the presence of clinical signs of HCM. Although no direct evidence comparing outcomes for at-risk individuals managed with and without genetic testing was identified, there is a strong chain of evidence that management changes can improve outcomes with genetic testing when there is a known familial variant. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who are asymptomatic with risk for HCM because of a positive family history who receive non-specific testing for an HCM-related variant, the evidence includes studies reporting on the clinical validity of testing. Relevant outcomes are overall survival, test accuracy and validity, changes in reproductive decision making, symptoms, and morbid events. Given the wide genetic variation in HCM and the likelihood that not all causative variants have been identified, there is imperfect clinical sensitivity. Therefore, a negative test is not sufficient to rule out a disease-associated variant in patients without a known family variant. For at-risk individuals without a known variant in the family, there is no clear relation between testing and improved outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Genetic testing for predisposition to hypertrophic cardiomyopathy (HCM) may be considered **medically necessary** for individuals who are at risk for development of HCM, defined as having a first-degree relative with established HCM, when there is a known pathogenic gene variant present in that affected relative (See Policy Guidelines).

Genetic testing for predisposition to HCM is considered **not medically necessary** for patients with a family history of HCM in which a first-degree relative with established HCM has tested negative for pathogenic variants.

Genetic testing for predisposition to HCM is considered **investigational** for all other patient populations, including but not limited to individuals who have a first-degree relative with clinical HCM, but in whom genetic testing is unavailable.

POLICY GUIDELINES

Due to the complexity of genetic testing for HCM and the potential for misinterpretation of results, the decision to test and the interpretation of test results should be performed by, or in consultation with, an expert in the area of medical genetics and/or HCM.

To inform and direct genetic testing for at-risk individuals, genetic testing should be initially performed in at least one close relative with definite HCM (index case), if possible. Recommendations indicate that, when possible, genetic testing for HCM be performed in an affected family member so that testing in unaffected, at-risk family members can focus on the mutation found in the affected family member. This testing is intended to document whether a known pathogenic variant is present in the family, and to optimize the predictive value of predisposition testing for at-risk relatives.

Because there are varying degrees of penetrance for different HCM mutations, consideration for testing of second- or third-degree relatives may be appropriate in certain circumstances. Some judgment should be allowed for these decisions (e.g., in the case of a small family pedigree). Consultation with an expert in medical genetics

and/or the genetics of HCM, in conjunction with a detailed pedigree analysis, is appropriate when testing of second- or third-degree relatives is considered.

GENETICS NOMENCLATURE UPDATE

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical protocol updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Genome Project, the HUMAN Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology—"pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign"—to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

GENETIC COUNSELING

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

MEDICARE ADVANTAGE

Genetic testing for inherited cardiomyopathy (e.g., hypertrophic cardiomyopathy, dilated cardiomyopathy, arrhythmogenic right ventricular cardiomyopathy) with a genomic sequence analysis panel is unlikely to impact therapeutic decision-making in the clinical management of the patient and is considered **not medically necessary**.

BACKGROUND**FAMILIAL HYPERTROPHIC CARDIOMYOPATHY**

Familial HCM is the most common genetic cardiovascular condition, with a phenotypic prevalence of approximately one (0.2%) in 500 adults.¹ It is the most common cause of sudden cardiac death (SCD) in adults younger than 35 years of age and is probably the most common cause of death in young athletes.² The overall mortality rate for patients with HCM is estimated to be 1% per year in the adult population.³

The genetic basis for HCM is a defect in the cardiac sarcomere, which is the basic contractile unit of cardiac myocytes and is composed of different protein structures.⁴ Around 1,400 disease-associated variants in at least 18 different genes have been identified.⁵⁻⁸ About 90% of pathogenic variants are missense (i.e., one amino acid is replaced for another), and the strongest evidence for pathogenicity is available for 11 genes coding for thick filament proteins (MYH7, MYL2, MYL3), thin filament proteins (TNNT2, TNNI3, TNNC1, TPM1, ACTC), intermediate filament proteins (MYBPC3), and the Z-disc adjoining the sarcomere (ACTN2, MYOZ2). Variants in myosin heavy chain (MYH7) and myosin-binding protein C (MYBPC3) are the most common and account for roughly 80% of sarcomeric HCM. These genetic defects are inherited in an autosomal dominant pattern with rare exceptions.⁴ In patients with clinically documented HCM, genetic abnormalities can be identified in approximately 60%.^{6,9} Most patients with the clinically documented disease are demonstrated to have a familial pattern, although some exceptions are found presumably due to de novo variants.⁹

Diagnosis and Management

The clinical diagnosis of HCM depends on the presence of left ventricular hypertrophy, measured by echocardiography or magnetic resonance imaging, in the absence of other known causative factors such as valvular disease, long-standing hypertension, or another myocardial disease.⁶ In addition to primary cardiac disorders, there are systemic diseases that can lead to left ventricular hypertrophy and thus mimic HCM. They include infiltrative diseases such as amyloidosis, glycogen storage diseases (e.g., Fabry disease, Pompe disease), and neuromuscular disorders (e.g., Noonan syndrome, Friedreich ataxia).⁹ These disorders need to be excluded before a diagnosis of familial HCM is made.

HCM is a very heterogeneous disorder. Manifestations range from subclinical, asymptomatic disease to severe, life-threatening disease. Wide phenotypic variability exists among individuals, even when an identical variant is present, including among affected family members.² This variability in clinical expression may be related to environmental factors and modifier genes.¹⁰ A large percentage of patients with HCM, perhaps the majority, are asymptomatic or have minimal symptoms.^{9,10} These patients do not require treatment and are not generally at high-risk for SCD. A subset of patients has severe disease that causes a major impact on quality of life and life expectancy. Severe disease can lead to disabling symptoms, as well as complications of HCM, including heart failure and malignant ventricular arrhythmias. Symptoms and presentation may include SCD due to unpredictable ventricular tachyarrhythmias, heart failure, or atrial fibrillation, or some combination.¹¹

Management of patients with HCM involves treating cardiac comorbidities, avoiding therapies that may worsen obstructive symptoms, treating obstructive symptoms with β -blockers, calcium channel blockers, and (if symptoms persist) invasive therapy with surgical myectomy or alcohol ablation, optimizing treatment for heart failure, if present, and SCD risk stratification. Implantable cardioverter defibrillator implantation may be indicated if there is a family history of SCD.

Diagnostic screening of first-degree relatives and other family members is an important component of HCM management. Guidelines have been established for screening clinically unaffected relatives of affected individuals. Screening with physical examination, electrocardiography, and echocardiography is recommended every 12 to 18 months for individuals ages 12 to 18 years and every three to five years for adults.¹⁰ Additional screening is recommended for any change in symptoms that might indicate the development of HCM.¹⁰

Genetic Testing

Genetic testing has been proposed as a component of screening at-risk individuals to determine predisposition to HCM among those patients at-risk. Patients at-risk for HCM are defined as individuals who have a close relative with established HCM. Results of genetic testing may influence the management of at-risk individuals, which may, in turn, lead to improved outcomes. Furthermore, results of genetic testing may have implications for decision making in the areas of reproduction, employment, and leisure activities. However, the likelihood of obtaining a positive genetic test in the proband is only about 50% because all genes causing HCM have not yet been identified or are absent from testing panels. Failure to identify the causative variant in the proband is an indeterminate result that provides no useful information and precludes predictive testing in 33% to 67% of cases.

Commercial testing has been available since 2003, and numerous companies offer genetic testing for HCM.^{5,12-15} Testing is performed either as a comprehensive or targeted gene test. Comprehensive testing, which is done for an individual without a known genetic variant in the family, analyzes the genes most commonly associated with genetic variants for HCM and evaluates whether any potentially pathogenic variants are present. Some available panels include testing for multisystem storage diseases that may include cardiac hypertrophy, such as Fabry disease (GLA), familial transthyretin amyloidosis (TTR), and X-linked Danon disease (LAMP2).

Other panels include testing for genes related to HCM and those associated with other cardiac disorders. For example, the Comprehensive Cardiomyopathy panel (ApolloGen) is a next-generation sequencing panel of 44 genes associated with HCM, dilated cardiomyopathy, restrictive cardiomyopathy, arrhythmogenic right ventricular cardiomyopathy, catecholaminergic polymorphic ventricular tachycardia, left ventricular noncompaction syndrome, Danon syndrome, Fabry disease, Barth syndrome, and transthyretin amyloidosis.¹⁶

For a patient with a known variant in the family, targeted testing is performed. Targeted variant testing evaluates for the presence or absence of a single variant known to exist in a close relative.

It can be difficult to determine the pathogenicity of genetic variants associated with HCM. Some studies have reported that assignment of pathogenicity has a relatively high error rate and that classification changes over time.^{17,18} With next-generation sequencing and whole-exome sequencing techniques, the sensitivity of identifying variants on the specified genes has increased substantially. At the same time, the number of variants of uncertain significance is also increased with next-generation sequencing. Also, the percentage of individuals who have more than one variant that is thought to be pathogenic is increasing. A 2013 study reported that 9.5% (19/200) patients from China with HCM had multiple pathogenic variants and that the number of variants correlated with severity of disease.¹⁹

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Sequencing tests for HCM are available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

No assay kits have been approved by the Food and Drug Administration for genetic testing for HCM.

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Semsarian C, Ingles J, Maron MS, et al. New perspectives on the prevalence of hypertrophic cardiomyopathy. *J Am Coll Cardiol*. Mar 31 2015;65(12):1249-1254. PMID 25814232
2. Alcalai R, Seidman JG, Seidman CE. Genetic basis of hypertrophic cardiomyopathy: from bench to the clinics. *J Cardiovasc Electrophysiol*. Jan 2008;19(1):104-110. PMID 17916152
3. Spirito P, Autore C, Formisano F, et al. Risk of sudden death and outcome in patients with hypertrophic cardiomyopathy with benign presentation and without risk factors. *Am J Cardiol*. May 1 2014;113(9):1550-1555. PMID 24630786
4. Keren A, Syrris P, McKenna WJ. Hypertrophic cardiomyopathy: the genetic determinants of clinical disease expression. *Nat Clin Pract Cardiovasc Med*. Mar 2008;5(3):158-168. PMID 18227814
5. Maron BJ, Maron MS, Semsarian C. Genetics of hypertrophic cardiomyopathy after 20 years: clinical perspectives. *J Am Coll Cardiol*. Aug 21 2012;60(8):705-715. PMID 22796258
6. Cirino AL, Ho C. Hypertrophic Cardiomyopathy Overview. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2014.
7. Ghosh N, Haddad H. Recent progress in the genetics of cardiomyopathy and its role in the clinical evaluation of patients with cardiomyopathy. *Curr Opin Cardiol*. Mar 2011;26(2):155-164. PMID 21297463
8. Teo LY, Moran RT, Tang WH. Evolving approaches to genetic evaluation of specific cardiomyopathies. *Curr Heart Fail Rep*. Dec 2015;12(6):339-349. PMID 26472190
9. Elliott P, McKenna WJ. Hypertrophic cardiomyopathy. *Lancet*. Jun 5 2004;363(9424):1881-1891. PMID 15183628
10. Maron BJ, McKenna WJ, Danielson GK, et al. American College of Cardiology/European Society of Cardiology clinical expert consensus document on hypertrophic cardiomyopathy. A report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents and the European Society of Cardiology Committee for Practice Guidelines. *J Am Coll Cardiol*. Nov 5 2003;42(9):1687-1713. PMID 14607462
11. Gersh BJ, Maron BJ, Bonow RO, et al. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. Nov 8 2011;124(24):2761-2796. PMID 22068434
12. Arya A, Bode K, Piorkowski C, et al. Catheter ablation of electrical storm due to monomorphic ventricular tachycardia in patients with nonischemic cardiomyopathy: acute results and its effect on long-term survival. *Pacing Clin Electrophysiol*. Dec 2010;33(12):1504-1509. PMID 20636312
13. GeneDx. Personal Communication. April 29, 2010.
14. Correlagan. Personal Communication. April 27, 2010.
15. PGxHealth. Personal Communication. April 22, 2010.
16. ApolloGen. Comprehensive Cardiomyopathy Panel. <https://www.ncbi.nlm.nih.gov/gtr/tests/525912/>. Accessed February 13, 2019.

17. Das KJ, Ingles J, Bagnall RD, et al. Determining pathogenicity of genetic variants in hypertrophic cardiomyopathy: importance of periodic reassessment. *Genet Med*. Apr 2014;16(4):286-293. PMID 24113344
18. Andreasen C, Nielsen JB, Refsgaard L, et al. New population-based exome data are questioning the pathogenicity of previously cardiomyopathy-associated genetic variants. *Eur J Hum Genet*. Sep 2013;21(9):918-928. PMID 23299917
19. Zou Y, Wang J, Liu X, et al. Multiple gene mutations, not the type of mutation, are the modifier of left ventricle hypertrophy in patients with hypertrophic cardiomyopathy. *Mol Biol Rep*. Jun 2013;40(6):3969-3976. PMID 23283745
20. BlueCross BlueShield Association Technology Evaluation Center (TEC). Genetic testing for predisposition to inherited hypertrophic cardiomyopathy. *TEC Assessment*. 2009;24(11).
21. Niimura H, Bachinski LL, Sangwatanaroj S, et al. Mutations in the gene for cardiac myosin-binding protein C and late-onset familial hypertrophic cardiomyopathy. *N Engl J Med*. Apr 30 1998;338(18):1248-1257. PMID 9562578
22. Richard P, Charron P, Carrier L, et al. Hypertrophic cardiomyopathy: distribution of disease genes, spectrum of mutations, and implications for a molecular diagnosis strategy. *Circulation*. May 6 2003;107(17):2227-2232. PMID 12707239
23. Van Driest SL, Ommen SR, Tajik AJ, et al. Sarcomeric genotyping in hypertrophic cardiomyopathy. *Mayo Clin Proc*. Apr 2005;80(4):463-469. PMID 15819282
24. Van Driest SL, Jaeger MA, Ommen SR, et al. Comprehensive analysis of the beta-myosin heavy chain gene in 389 unrelated patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. Aug 4 2004;44(3):602-610. PMID 15358028
25. Van Driest SL, Ellsworth EG, Ommen SR, et al. Prevalence and spectrum of thin filament mutations in an outpatient referral population with hypertrophic cardiomyopathy. *Circulation*. Jul 29 2003;108(4):445-451. PMID 12860912
26. Mogensen J, Murphy RT, Kubo T, et al. Frequency and clinical expression of cardiac troponin I mutations in 748 consecutive families with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. Dec 21 2004;44(12):2315-2325. PMID 15607392
27. Michels M, Soliman OI, Phefferkorn J, et al. Disease penetrance and risk stratification for sudden cardiac death in asymptomatic hypertrophic cardiomyopathy mutation carriers. *Eur Heart J*. Nov 2009;30(21):2593-2598. PMID 19666645
28. Cardoso B, Gomes I, Loureiro P, et al. Clinical and genetic diagnosis of familial hypertrophic cardiomyopathy: Results in pediatric cardiology. *Rev Port Cardiol*. Mar 2017;36(3):155-165. PMID 28214152
29. van Velzen, HH, Schinkel, AA, Baart, SS, Oldenburg, RR, Frohn-Mulder, II, van Slegtenhorst, MM, Michels, MM. Outcomes of Contemporary Family Screening in Hypertrophic Cardiomyopathy. *Circ Genom Precis Med*, 2018 Apr 18;11(4). PMID 29661763
30. Ko, CC, Arscott, PP, Concannon, MM, Saberi, SS, Day, SS, Yashar, BB, Helms, AA. Genetic testing impacts the utility of prospective familial screening in hypertrophic cardiomyopathy through identification of a nonfamilial subgroup. *Genet. Med.*, 2017 Jun 24;20(1). PMID 28640247
31. Lopes, LL, Brito, DD, Belo, AA, Cardim, NN. Genetic characterization and genotype-phenotype associations in a large cohort of patients with hypertrophic cardiomyopathy - An ancillary study of the Portuguese registry of hypertrophic cardiomyopathy. *Int. J. Cardiol.*, 2018 Dec 18;278:173-179. PMID 30554928
32. Sedaghat-Hamedani, FF, Kayvanpour, EE, Tugrul, OO, Lai, AA, Amr, AA, Haas, JJ, Proctor, TT, Ehlermann, PP, Jensen, KK, Katus, HH, Meder, BB. Clinical outcomes associated with sarcomere mutations in hypertrophic cardiomyopathy: a meta-analysis on 7675 individuals. *Clin Res Cardiol*, 2017 Aug 26;107(1). PMID 28840316
33. Erdmann J, Daehmlow S, Wischke S, et al. Mutation spectrum in a large cohort of unrelated consecutive patients with hypertrophic cardiomyopathy. *Clin Genet*. Oct 2003;64(4):339-349. PMID 12974739
34. Olivotto I, Girolami F, Ackerman MJ, et al. Myofilament protein gene mutation screening and outcome of patients with hypertrophic cardiomyopathy. *Mayo Clin Proc*. Jun 2008;83(6):630-638. PMID 18533079

35. Chiou KR, Chu CT, Charng MJ. Detection of mutations in symptomatic patients with hypertrophic cardiomyopathy in Taiwan. *J Cardiol*. Mar 2015;65(3):250-256. PMID 25086479
36. Adalsteinsdottir B, Teekakirikul P, Maron BJ, et al. Nationwide Study on Hypertrophic Cardiomyopathy in Iceland: Evidence of a MYBPC3 Founder Mutation. *Circulation*. Sep 30 2014;130(14):1158-1167. PMID 25078086
37. Ingles, JJ, Goldstein, JJ, Thaxton, CC, Caleshu, CC, Corty, EE, Crowley, SS, Dougherty, KK, Harrison, SS, McGlaughon, JJ, Milko, LL, Morales, AA, Seifert, BB, Strande, NN, Thomson, KK, van Tintelen, JJ, Wallace, KK, Walsh, RR, Wells, QQ, Whiffin, NN, Witkowski, LL, Semsarian, CC, Ware, JJ, Hershberger, RR, Funke, BB. Evaluating the Clinical Validity of Hypertrophic Cardiomyopathy Genes. *Circ Genom Precis Med*, 2019 Jan 27. PMID 30681346
38. Authors/Task Force m, Elliott PM, Anastasakis A, et al. 2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: The Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). *Eur Heart J*. Oct 14 2014;35(39):2733-2779. PMID 25173338
39. Maron, BB, Zipes, DD, Kovacs, RR. Eligibility and Disqualification Recommendations for Competitive Athletes With Cardiovascular Abnormalities: Preamble, Principles, and General Considerations: A Scientific Statement From the American Heart Association and American College of Cardiology. *Circulation*, 2015 Dec 2;132(22). PMID 26621642
40. Maron, BB, Udelson, JJ, Bonow, RR, Nishimura, RR, Ackerman, MM, Estes, NN, Cooper, LL, Link, MM, Maron, MM. Eligibility and Disqualification Recommendations for Competitive Athletes With Cardiovascular Abnormalities: Task Force 3: Hypertrophic Cardiomyopathy, Arrhythmogenic Right Ventricular Cardiomyopathy and Other Cardiomyopathies, and Myocarditis: A Scientific Statement From the American Heart Association and American College of Cardiology. *J. Am. Coll. Cardiol.*, 2015 Nov 7;66(21). PMID 26542657
41. Ackerman MJ, Priori SG, Willems S, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm*. Aug 2011;8(8):1308-1339. PMID 21787999
42. National Government Services, Inc. (Primary Geographic Jurisdiction 06 & K- Illinois, Minnesota, Wisconsin, Connecticut, New York - Entire State, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont) Local Coverage Article: Billing and Coding: Molecular Pathology Procedures (A56199), Revision Effective Date 10/15/2019.