

Protocol

Genetic Testing for Breast Cancer Gene Expression Prognosis Assay

(20436, 20476)

Medical Benefit		Effective Date: 01/01/19	Next Review Date: 09/19
Preauthorization	Yes	Review Dates: 03/08, 03/09, 01/10, 01/11, 01/12, 01/13, 01/14, 11/14, 11/15, 09/16, 03/17, 09/17, 09/18	

Preauthorization is required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: • With early-stage node-negative invasive breast cancer considering adjuvant chemotherapy	Interventions of interest are: • Gene expression profiling with Oncotype DX (21-gene signature), Endopredict, Breast Cancer Index, MammaPrint (70-gene signature), Prosigna	Comparators of interest are: • Clinical risk prediction algorithms	Relevant outcomes include: • Disease-specific survival • Change in disease status
Individuals: • With early-stage node-positive invasive breast cancer considering adjuvant chemotherapy	Interventions of interest are: • Gene expression profiling with Oncotype DX (21-gene signature), Endopredict, MammaPrint (70-gene signature), Prosigna	Comparators of interest are: • Clinical risk prediction algorithms	Relevant outcomes include: • Disease-specific survival • Change in disease status
Individuals: • With ductal carcinoma in situ considering radiotherapy	Interventions of interest are: • Gene expression profiling with the Oncotype DX Breast DCIS Score	Comparators of interest are: • Clinical risk prediction algorithms	Relevant outcomes include: • Change in disease status
Individuals: • With early-stage node-negative invasive breast cancer recurrence-free at five years, considering extended endocrine therapy	Interventions of interest are: • Gene expression profiling with Oncotype DX (21-gene signature), EndoPredict, Breast Cancer Index, MammaPrint (70-gene signature) or Prosigna	Comparators of interest are: • Clinical risk prediction algorithms	Relevant outcomes include: • Disease-specific survival • Change in disease status
Individuals: • With breast cancer who are undergoing assessment of HER2 status	Interventions of interest are: • Quantitative total HER2 protein expression and HER2 homodimer measurement	Comparators of interest are: • Assessment of HER2 status using immunohistochemistry or fluorescence in situ hybridization	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity

DESCRIPTION

Laboratory tests have been developed that detect the expression, via messenger RNA, of many different genes in breast tumor tissue and combine the results into a prediction of distant recurrence risk for women with early-stage breast cancer. Test results may help providers and patients decide whether to include adjuvant chemotherapy in the postsurgical management of breast cancer to alter treatment in patients with ductal carcinoma in situ (DCIS) or to recommend extended endocrine therapy in patients who are recurrence-free at five years).

Novel assays that quantitatively measure total human epidermal growth factor receptor 2 (HER2) protein expression and homodimers have been developed to improve the accuracy and consistency of HER2 testing.

SUMMARY OF EVIDENCE

For all tests and all indications, relevant outcomes include disease-specific survival and changes in disease status.

EARLY-STAGE NODE-NEGATIVE INVASIVE BREAST CANCER

For the evaluation of breast cancer-related gene expression profiling tests for the management of all early-stage breast cancer populations, study populations considered had positive hormone receptor status, and negative human epidermal growth factor receptor 2 (HER2) status. Studies retrospectively collecting tumor samples from prospective trials that provide 10-year distant recurrence rates or 10-year survival rates in node-negative women not receiving adjuvant chemotherapy were included.

Oncotype DX (21-Gene Assay)

For individuals who have early-stage node-negative invasive breast cancer considering adjuvant chemotherapy who receive gene expression profiling with Oncotype DX (21-gene assay), the evidence includes multiple prospective clinical trials and prospective-retrospective studies. Patients classified as low risk with Oncotype DX have a low risk of recurrence in which avoidance of adjuvant chemotherapy is reasonable (average risk at 10 years, 3%-7%; upper bound of the 95% confidence interval (CI), 6% to 10%). These results have been demonstrated with stronger study designs for evaluating biomarkers. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

EndoPredict

For individuals who have early-stage node-negative invasive breast cancer considering adjuvant chemotherapy who receive gene expression profiling with EndoPredict, the evidence includes three prospective-retrospective studies and observational studies. The studies revealed that a low score was associated with a low absolute risk of 10-year distant recurrence (average risk at 10 years for the two larger studies, 3%-6%; upper bound of the 95% CI, 6% to 9%). Over half of patients in these studies were classified at low risk. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Prosigna

For individuals who have early-stage node-negative invasive breast cancer considering adjuvant chemotherapy who receive gene expression profiling with Prosigna, the evidence includes two prospective-retrospective studies evaluating the prognostic ability of Prosigna. Both studies showed a low absolute risk of distant recurrence in patients with low risk scores (average risk at 10 years, 3%-5%; upper bound for the study providing CI, 6%). The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

EARLY-STAGE NODE-POSITIVE INVASIVE BREAST CANCER

Oncotype DX (21-Gene Assay)

For individuals who have early-stage node-positive invasive breast cancer who are considering adjuvant chemotherapy who receive gene expression profiling with Oncotype DX (21-gene assay), the evidence includes two prospective-retrospective studies and a prospective study. The prospective-retrospective studies showed that Oncotype DX stratifies node-positive patients into high and low risk for distant recurrence-free survival. However, only one of the studies reported CIs for estimates and those are very wide. The prospective study included patients with node-negative and node-positive breast cancer. The authors reported that subgroup analyses of patients with node-positive breast cancer who were classified as low risk experienced higher rates of survival than patients classified as high risk, though no rates were provided. There is a wide range of survival improvements over which individual patients would elect or refuse adjuvant chemotherapy, but accurate risk estimates are needed to inform patient decisions. The evidence is insufficient to determine the effects of the technology on health outcomes.

EndoPredict

For individuals who have early-stage node-positive invasive breast cancer who are considering adjuvant chemotherapy who receive gene expression profiling with EndoPredict, the evidence includes two prospective-retrospective analyses. In a study, the 10-year distant recurrence rate in low-risk EPclin score patients was estimated to be 5% (95% CI, 1% to 9%). In the other study, 10-year distant recurrence rate in low-risk EPclin score patients was estimated to be 5%, but the upper bound of the 95% CI was close to 20%. To establish that the test has potential for clinical utility, it should be able to identify a low-risk group with a recurrence risk that falls within a range that is clinically meaningful for decision-making about avoiding adjuvant chemotherapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

MammaPrint (70-Gene Signature)

For individuals who have early-stage node-positive invasive breast cancer who are considering adjuvant chemotherapy who receive gene expression profiling with MammaPrint (70-gene signature), the evidence includes a clinical utility study and an observational study. The study of clinical utility only reported five-year results and may not identify a group with sufficiently low risk. The observational study reported that the low-risk group experienced a low rate of 10-year distant recurrence; however, the standard error around the rate did not meet the threshold benefit of less than 10%. The evidence is insufficient to determine the effects of the technology on health outcomes.

Prosigna

For individuals who have early-stage node-positive invasive breast cancer who are considering adjuvant chemotherapy who receive gene expression profiling with the Prosigna risk of recurrence (ROR) score, the evidence includes a single prospective-retrospective study. The 10-year distant recurrence rate in low-risk Prosigna ROR patients with a single positive node is roughly twofold the rate in low-risk ROR score node-negative patients. However, in the single available study, the upper bound of the 95% CI for 10-year distant recurrence in node-positive patients classified as ROR score low-risk was about 13%, which approaches the range judged clinically informative in node-negative patients. The predicted recurrence rates require replication. To establish that the test has potential for clinical utility, it should be able to identify a low-risk group with a recurrence risk that falls within a range that is clinically meaningful for decision-making about avoiding adjuvant chemotherapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

Extended Endocrine Therapy

For this indication, Oncotype DX, EndoPredict, Breast Cancer Index, MammaPrint, and Prosigna were evaluated. Studies retrospectively collecting tumor samples from prospective trials that provided 10-year distant recur-

rence rates or 10-year survival rates were included. Studies comparing genetic assays with clinical risk prediction tools were also included.

Oncotype DX (21-Gene Assay)

For individuals who have early-stage node-negative invasive breast cancer who are distant recurrence-free at five years who are considering extending endocrine treatment who receive gene expression profiling with Oncotype DX (21-gene assay), the evidence includes a study from a previously conducted clinical trial. The study did not show low distant recurrence rates in patients classified as low risk with the test, and no CIs were presented. The ability of the test to reclassify patients assessed with a clinical prediction tool was not reported. The evidence is insufficient to determine the effects of the technology on health outcomes.

EndoPredict

For individuals who have early-stage node-negative invasive breast cancer who are distant recurrence-free at five years who are considering extending endocrine treatment who receive gene expression profiling with EndoPredict, the evidence includes a study of archived tissue samples from a previously conducted clinical trial. The study showed low distant recurrence rates in patients classified as low risk with EndoPredict. The ability of the test to reclassify patients assessed with a clinical prediction tool was not reported. Additional prospective trials or retrospective-prospective studies of archived samples reporting on the association between risk score and survival are needed for confirmation of results from the single study. More importantly, clarity is needed about how the test would inform clinical practice. The evidence is insufficient to determine the effects of the technology on health outcomes.

Breast Cancer Index

For individuals who have early-stage node-negative invasive breast cancer who are distant recurrence-free at five years who are considering extending tamoxifen treatment who receive gene expression profiling with the Breast Cancer Index, the evidence includes two studies of archived tissue samples from previously conducted clinical trials and a retrospective cohort study. The three studies showed low distant recurrence rates in patients classified as low risk with the test. Two studies suggested that, in addition to having a more favorable prognosis, low-risk patients may receive lesser benefit from extended endocrine therapy. The ability of the test to reclassify patients assessed with a clinical prediction tool was not reported. Clarity about how the test would inform clinical practice is needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

MammaPrint (70-Gene Signature)

For individuals who have early-stage node-negative invasive breast cancer who are distant recurrence-free at five years who are considering extending tamoxifen treatment who receive gene expression profiling with MammaPrint (70-gene signature), the evidence includes a retrospective-prospective study. Analyses on patients classified as ultralow risk (a subgroup of the low-risk group) showed that this ultralow-risk group experienced high 10- and 20-year breast cancer-specific survival rates. Additional studies are needed to confirm the results of this single study. The ability of the test to reclassify patients assessed with a clinical prediction tool was not reported. Clarity about how the test would inform clinical practice is needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Prosigna

For individuals who have early-stage node-negative invasive breast cancer who are distant recurrence-free at five years who are considering extending tamoxifen treatment who receive gene expression profiling with Prosigna, the evidence includes two studies from previously conducted clinical trials examined in three publications. The studies showed low distant recurrence rates in patients classified as low risk with the test. A reclassification result suggested that the test may offer little improvement over clinical predictors alone. Clarity about

how the test would inform clinical practice is needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

HER2

For individuals who have breast cancer and are undergoing assessment of HER2 status who receive quantitative total HER2 protein expression and HER2 homodimer measurement, the evidence includes validation studies and retrospective analysis of the association between levels and survival outcomes. Relevant outcomes are overall survival, disease-specific survival, test accuracy, and test validity. Retrospective analysis using HERmark have shown that the assay may predict a worse response to trastuzumab in certain populations. However, findings have been inconsistent, and no clear association with clinical outcomes has been shown. Additionally, cut points for defining patient groups varied across studies. Clinical utility of the HERmark assay has not been demonstrated, and clinical trials are needed to determine the impact on clinical outcomes of patients stratified by the HERmark assay. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

The use of gene expression assays for deciding the value of adjuvant therapy in individuals with stage I breast cancer may be considered **medically necessary** for women meeting all of the following characteristics:

- Unilateral tumor; AND
- hormone receptor-positive (that is estrogen-receptor [ER]-positive or progesterone receptor [PR]-positive); AND
- human epidermal growth factor receptor 2 (HER2) negative; AND
- tumor size greater than 0.5 cm with moderate/poor differentiation or unfavorable features; AND
- node negative or lymph nodes with micrometastases less than two mm in size; AND
- who will be treated with adjuvant endocrine therapy, e.g., tamoxifen or aromatase inhibitors; AND
- when the test result will aid the patient in making the decision regarding chemotherapy (i.e., when chemotherapy is a therapeutic option); AND
- when ordered within six months after diagnosis.

The use of the Oncotype DX, Prosigna or Endopredict RT-PCR, or the MammaPrint® assays may be considered **medically necessary** if the individual meets the criteria above.

In addition, for individuals who have stage II breast cancer, who are at a high clinical risk of recurrence and who meet all other indications as above, but who have one to three positive lymph nodes (in place of “node negative or lymph nodes with micrometastases less than two mm in size”), the use of MammaPrint to inform decisions on adjuvant systemic chemotherapy may be considered **medically necessary**.

The use of MammaPrint is considered **investigational** in individuals who have a low clinical risk for recurrence (per MINDACT trial categorization.)

The 21-gene RT-PCR assay Oncotype DX™ should only be ordered on a tissue specimen obtained during surgical removal of the tumor and after subsequent pathology examination of the tumor has been completed and determined to meet the above criteria (i.e., the test should not be ordered on a preliminary core biopsy). The test should be ordered in the context of a physician-patient discussion regarding risk preferences and when the test result will aid the patient in making decisions regarding chemotherapy.

For patients who otherwise meet the above characteristics but who have multiple ipsilateral primary tumors, a specimen from the tumor with the most aggressive histological characteristics should be submitted for testing. It is not necessary to conduct testing on each tumor; treatment is based on the most aggressive lesion.

All other indications for multigene breast cancer panel assays, including determination of recurrence risk in invasive breast cancer patients with positive lymph nodes, patients with bilateral disease, or to consider length of treatment with tamoxifen are considered **investigational**.

The use of a subset of genes from the Oncotype DX or Prosigna RT-PCR assay for predicting recurrence risk in individuals with noninvasive ductal carcinoma in situ (i.e., Oncotype DX® Breast DCIS Score) to inform treatment planning following excisional surgery is considered **investigational**.

The use of other gene expression assays, including but not limited to, Mammostrat® Breast Cancer Test, the Breast Cancer IndexSM, BreastOncPxTM, NexCourse® Breast IHC4 and BreastPRSTM, for any indication is considered **investigational**.

The use of gene expression assays in men with breast cancer is considered **investigational**.

The use of gene expression assays to molecularly subclassify breast cancer is considered **investigational**.

The use of gene expression assays for quantitative assessment of ER, PR, and HER2 overexpression is considered **investigational**.

POLICY GUIDELINES

The NCCN Panel members acknowledge that many assays have been clinically validated for prediction of prognosis. However, based on the currently available data, the panel believes that the 21-gene assay has been best validated for its use as a prognostic test as well as in predicting who is most likely to respond to systemic chemotherapy.

*Women meeting these criteria whose MammaPrint score is low may be treated with hormone therapy alone, as it is unlikely that chemotherapy will provide substantial additional benefits (per MINDACT trial categorization).

GENETIC COUNSELING

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

MEDICARE ADVANTAGE

The above medical necessity criteria applies for Medicare Advantage members for all gene expression assays in breast tumor tissue tests when no separate Medicare Advantage criteria exists.

PROSIGNA® Breast Cancer Prognostic Gene Signature Assay is considered **medically necessary** in patients who have undergone surgery in conjunction with locoregional treatment consistent with standard of care, either as:

- A prognostic indicator for distant recurrence-free survival at 10 years in post-menopausal women with Hormone Receptor-Positive (HR+), lymph node-negative, Stage I or II breast cancer to be treated with adjuvant endocrine therapy alone, when used in conjunction with other clinicopathological factors.
- A prognostic indicator for distant recurrence-free survival at 10 years in post-menopausal women with Hormone Receptor-Positive (HR+), lymph node-positive (one to three positive nodes), Stage II breast cancer to be treated with adjuvant endocrine therapy alone, when used in conjunction with other clinicopathological factors. The device is not intended for patients with four or more positive nodes.

The Breast Cancer Index (aka BCI) (bioTheranostics) may be considered **medically necessary** for members that meet the following criteria:

- Post-menopausal female with non-relapsed, ER+ BREAST CANCER, and
- Is lymph node negative, and
- Is completing five (5) years of tamoxifen therapy, and
- Patient must be eligible for consideration of extended endocrine therapy based on published clinical trial data or practice guidelines, and
- Physician or patient is concerned about continuing anti-hormonal therapy because of documented meaningful toxicity or possible significant patient-specific side effects, and
- The test results will be discussed with the patient (including the limitations of the testing method, the risks and benefits of either continuing or stopping the therapy based on the test, and current cancer management guidelines).

Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 21 genes (Oncotype DX), utilizing formalin-fixed paraffin embedded tissue, algorithm reported as recurrence score is considered **medically necessary** to guide therapeutic decision-making in patients with the following findings:

- estrogen-receptor positive, node-negative carcinoma of the breast
- estrogen-receptor positive micrometastases of carcinoma of the breast, and
- estrogen-receptor positive breast carcinoma with one to three positive nodes.

For Medicare Advantage the HERmark®, and the MammaPrint™ tests are considered **medically necessary**.

For Medicare Advantage the Oncotype DX® DCIS assay (Genomic Health, Inc., Redwood City, CA) is considered **medically necessary** for women diagnosed with DCIS who are planning on having breast conserving surgery and considering adjuvant radiation therapy only when the following clinical conditions are met:

- Pathology (excisional or core biopsy) reveals ductal carcinoma in situ of the BREAST (no pathological evidence of invasive disease), and
- FFPE specimen with at least 0.5 mm of DCIS length, and
- Patient is a candidate for and is considering breast conserving surgery alone as well as breast conserving surgery combined with adjuvant radiation therapy, and
- Test result will be used to determine treatment choice between surgery alone vs. surgery with radiation therapy, and
- Patient has not received and is not planning on receiving a mastectomy.

For Medicare Advantage, the EndoPredict® breast cancer gene expression test (Myriad Genetic Laboratories Inc., Salt Lake City, UT) is considered **medically necessary** for the management of post-menopausal women

diagnosed with early-stage (TNM stage T1-3, N0-1) estrogen-receptor (ER) positive, Her2-negative breast cancer, who are either lymph node-negative or who have 1-3 positive nodes, and for whom treatment with adjuvant endocrine therapy (e.g., tamoxifen or aromatase inhibitors) is being considered.

BACKGROUND

NEWLY DIAGNOSED BREAST CANCER

Most women with newly diagnosed breast cancer in the United States present with early-stage or locally advanced (i.e., nonmetastatic) disease. However, almost a third of women who are disease-free after initial local and regional treatment develop distant recurrences during follow-up.¹ Current breast cancer treatment regimens involve systemic adjuvant chemotherapy, hormonal therapy, biologic therapy, or a combination, depending on patients' baseline level of recurrence risk, hormonal markers, and risk tolerance.

Women whose tumors are positive for human epidermal growth factor receptor 2 (HER2) should receive adjuvant therapy with a HER2-directed therapy (trastuzumab with or without pertuzumab). Decision making about adjuvant biologic therapy for women with HER2-positive cancer is not discussed here. This review focuses on three decision points:

1. The decision to pursue adjuvant chemotherapy following locoregional therapy, with or without neoadjuvant chemotherapy, based on the predicted risk of recurrence, for women who are hormone receptor-positive but HER2-negative. The use of adjuvant chemotherapy reduces the risk of breast cancer recurrence but carries risks of systemic toxicity. The risk:benefit ratio must be balanced for each patient, with a higher likelihood of net health benefits for patients with a greater baseline predicted the risk of recurrence. Some of the individual considerations are discussed below. HER2 expression independently confers an unfavorable prognosis, but assessing the independent effects of HER2 is complicated in the presence of targeted therapy; therefore, we focus specifically on patients without HER2 expression.
2. The decision to pursue adjuvant endocrine therapy from five to 10 years for women who are hormone receptor-positive but HER2-negative and who have survived without recurrence for five years. For patients with hormone receptor-positive tumors, the use of adjuvant endocrine therapy (tamoxifen and/or an aromatase inhibitor, with or without ovarian suppression) for five to 10 years after an initial diagnosis has support in clinical practice. The 2017 guidelines from the National Comprehensive Cancer Network recommend extended endocrine therapy.² The American Society for Clinical Oncology's 2014 focused update to its guidelines on adjuvant endocrine therapy for women with hormone receptor-positive breast cancer have recommended 10 years of tamoxifen for pre- or perimenopausal women, and a total of seven to ten years of endocrine therapy, following one of four regimens that include tamoxifen with or without an aromatase inhibitor for postmenopausal women.^{3,4}
3. The decision to pursue adjuvant radiotherapy in women with ductal carcinoma in situ (DCIS). Adjuvant radiotherapy reduces the risk of local recurrences but has not been shown to change the risk of distant recurrence or mortality. There may be a group of patients for whom the reduction in risk for local recurrence may not be large enough to justify the risks of radiotherapy.

SELECTION OF ADJUVANT CHEMOTHERAPY BASED ON RISK OF RECURRENCE

An important part of treatment planning for women with breast cancer involves determining which patients could benefit from adjuvant cytotoxic chemotherapy. For example, for women with early-stage invasive breast cancer (i.e., cancer extending beyond the basement membrane of the mammary ducts into adjacent tissue), adjuvant cytotoxic chemotherapy consistently provides approximately a 30% relative risk reduction in 10-year breast cancer mortality regardless of patients' baseline prognosis. However, the absolute benefit of chemotherapy depends on the underlying or baseline risk of recurrence. Women with the best prognosis have tumors that

are small, early-stage, estrogen receptor–positive, and lymph node negative (Table 1 shows recurrence risk for estrogen receptor–positive cancers for patients followed in the International Breast Cancer Study Group).¹ Patients may have received no adjuvant treatment, or adjuvant tamoxifen and/or adjuvant chemotherapy. These women have an approximately 15% 10-year risk of recurrence with tamoxifen alone; approximately 85% of these patients could avoid the toxicity of adjuvant cytotoxic chemotherapy if they could be accurately identified. Conventional risk classifiers (e.g., Adjuvant! Online) estimate recurrence risk by considering criteria such as tumor size, type, grade, and histologic characteristics; hormone receptor status; and number of affected lymph nodes. Consensus guidelines for defining receptor status exist⁵; however, no single classifier is considered a criterion standard. As a result, a substantial number of patients are treated with chemotherapy who fail to benefit. Better predictors of recurrence risk could help women’s decision-making, some of whom may prefer to avoid chemotherapy if assured their risk is low.

Table 1. Effect of Nodal Involvement, Tumor Size, and Grade on Annual Recurrence Hazard in Estrogen Receptor-Positive Breast Cancers

Nodes	Recurrence, Hazard ^a (SE), %				
	Years				
	0-5	5-10	10-15	15-20	20-25
0	5.8 (0.5)	3.3 (0.4)	2.0 (0.4)	2.1 (0.4)	1.1 (0.4)
1 to 3	9.5 (0.6)	5.8 (0.6)	3.0 (0.5)	3.5 (0.7)	1.5 (0.6)
≥ 4	17.2 (0.9)	10.9 (1.2)	5.9 (1.2)	3.8 (1.2)	1.3 (0.9)
Size					
≤ 2 cm	7.0 (0.4)	4.8 (0.4)	2.9 (0.4)	2.7 (0.5)	1.5 (0.5)
> 2 cm	12.9 (0.6)	6.1 (0.6)	2.9 (0.5)	2.7 (0.5)	1.1 (0.5)
Grade					
1	5.8 (0.6)	4.9 (0.7)	3.6 (0.7)	4.0 (0.9)	0.7 (0.5)
2	9.6 (0.5)	6.3 (0.5)	2.8 (0.4)	2.7 (0.5)	1.8 (0.5)
3	14.1 (0.8)	4.1 (0.6)	2.5 (0.6)	2.4 (0.7)	0.4 (0.4)

Adapted from Colleoni et al (2016).¹

^a Number of events occurring within a time interval divided by the total years of follow-up during the interval accrued by patients at risk during the interval. Patients may have received no adjuvant treatment or have been treated with adjuvant tamoxifen and/or adjuvant chemotherapy.

SELECTION OF EXTENDED ENDOCRINE THERAPY

Randomized controlled trials (RCTs) have established that five years of tamoxifen improves mortality in women with hormone receptor-positive breast cancer. A 2011 individual patient data meta-analysis by the Early Breast Cancer Trialists’ Collaborative Group, including 20 trials (total N=21,457 patients) found that five years of tamoxifen in estrogen receptor–positive disease reduced the risk of recurrences by almost 50% over 10 years on the relative scale; breast cancer mortality was decreased by 29% through 15 years.⁶

For patients with early-stage, invasive breast cancer that is hormone receptor-positive, the use of endocrine therapy (tamoxifen and/or an aromatase inhibitor, with or without ovarian suppression) for five to 10 years following initial diagnosis has support in national guidelines.^{2, 3, 7} However, the regimens available and the evidence to support them vary.

RCTs published recently have shown that extended endocrine therapy decreases the risk of recurrence. The American Society for Clinical Oncology and the National Comprehensive Cancer Network guidelines were informed primarily by results of the ATLAS trial, which compared five and 10 years of tamoxifen⁸ and the subsequent aTTom trial (reported in abstract form).⁹ In both trials, in women who were hormone receptor-positive and had completed five years of tamoxifen, five years of extended tamoxifen was associated with improvements in breast cancer-specific mortality; ATLAS showed improvements in overall survival (see Table 2).

Three previously reported randomized trials of extended tamoxifen treatment had mixed findings: Tormey et al (1996; total N=194 patients),¹⁰ the National Surgical Adjuvant Breast and Bowel Project (Fisher et al, 2001; total N=1172 patients),¹¹ and the Scottish Cancer Trials Breast Group (Stewart et al, 2001; total N=342 patients)¹² (see Table 2).

Overall, the available trial evidence would suggest that 10 years of tamoxifen in pre- or postmenopausal women can be linked with improved survival while trials of extended aromatase inhibitors in different populations of hormone receptor-positive patients have had more mixed results.

Table 2. Randomized Trials Evaluating Adjuvant Extended Endocrine Therapies for Hormone Receptor-Positive Breast Cancer

Study	Population	Comparators	Breast Cancer–Specific Mortality		Overall Mortality	
			Event RR (95% CI)	p	Event RR (95% CI)	p
Extended tamoxifen						
ATLAS (2013) ⁸	6846 women with ER-positive, early breast cancer, after five years of tamoxifen	Continue tamoxifen to 10 years (n=3428) vs. stop tamoxifen at five years (n=3418)	0.83 (0.72 to 0.96) (331/3428 vs. 397/3418)	0.01	0.87 (0.78 to 0.97) 722 (639/3428 vs. 722/3418)	0.01
aTTom (2013) ⁹	6953 women with ER-positive or untested breast cancer, after five years of tamoxifen	Continue tamoxifen to 10 years (n=3468) vs. stop tamoxifen at five years (n=3485)	10 years 392/3468 intervention vs. 442/3485 control Years five-nine 1.03 (0.84 to 1.27) After year nine 0.77 (0.64 to 0.92)	0.05	10 years 849/3468 intervention vs. 910/3485 control Years five-nine 1.05 (0.90 to 1.22) After year nine 0.86 (0.75 to 0.97)	0.1
Extended aromatase inhibitor						
ABCSG (2007) ¹³	856 post-menopausal women with ER- and/or PR-positive breast cancer, after five years of tamoxifen	Anastrozole for three years (n=386) vs. no further therapy (n=466)			five years 10.3% anastrozole vs. 11.7% control Event HR (95% CI) 0.89 (0.59 to 1.34)	0.57
			Breast Cancer–Specific Survival		Overall Survival	
NCIC CTG MA.17 trial (2003, 2005) ^{14, 15}	5187 post-menopausal women with ER- and/or PR-positive early breast cancer, after five years tamoxifen	Continue letrozole to 10 years (n=2593) vs. stop tamoxifen at five years (n=2594)	48 Months 94.4% letrozole vs. 89.8% placebo Event HR 0.58 (0.45 to 0.76)	< 0.001	48 Months 96% letrozole vs. 94% placebo Event HR 0.76 (0.48 to 0.21) 40 Months 95.4% letrozole vs. 95% placebo Event HR 0.82 (0.57 to 1.19)	0.25 0.3
NSABP (2008) ¹⁶	1598 post-menopausal women with ER- and/or PR-positive early breast cancer, after five years of tamoxifen	Planned comparison: five years exemestane vs. five years placebo. Accrual stopped (n=1598 randomized) and crossover allowed after results of NCIC CTG available: • Exemestane: 783 randomized, 560 continued after	48 Months ITT: 91% exemestane vs. 89% placebo	0.07		

Study	Population	Comparators	Breast Cancer–Specific Mortality	Overall Mortality
		unblinding) • Placebo: 779 randomized, 334 crossed over to exemestane after unblinding		

ABCSG: Austrian Breast and Colorectal Cancer Study Group; CI: confidence interval; ER: estrogen receptor; HR: hazard ratio; ITT: intention to treat; NCIC CTG: National Cancer Institute Clinical Trials Group; NSABP: National Surgical Adjuvant Breast and Bowel Project; PR: progesterone receptor; RR: rate ratio.

In addition to the trials published in full-length form, three trials were presented in early 2017 evaluating extended endocrine therapy in postmenopausal women (NSABP-42 [NCT00382070]: 10 years vs. five years of letrozole; DATA [NCT00301457]: six years vs. three years of anastrozole; and IDEAL [NTR3077] 10 years vs. 7.5 years of letrozole) did not meet their primary end points.

CLINICAL USES OF GENE EXPRESSION SIGNATURES FOR BREAST CANCER

In other clinical scenarios involving breast cancer, accurate assessment of prognosis may affect the decision to offer certain treatments. Recently, several groups have identified panels of gene expression markers (“signatures”) that appear to predict the baseline risk of invasive breast cancer recurrence after surgery, radiotherapy, and endocrine therapy (for hormone receptor-positive tumors). Several gene expression tests commercially available in the United States are listed in Table 3. If these panels are more accurate risk predictors than current conventional classifiers, they could be used to aid decision making on adjuvant treatments without greatly affecting disease-free survival and overall survival. This review focuses on gene expression profiling panels that have prognostic or predictive ability in individuals with early-stage, invasive breast cancer with known estrogen receptor and progesterone receptor and HER2 status. The proposed clinical utility of these tests varies by the clinical context; these specific indications are discussed in this review:

1. Prognosis and/or prediction of treatment response in patients with node-negative, early-stage, hormone receptor-positive, HER2-negative invasive breast cancer who will receive adjuvant hormonal therapy for the purpose of determining whether patients can avoid adjuvant cytotoxic chemotherapy.
2. Prognosis and/or prediction of treatment response in patients with node-positive (one to three nodes), hormone receptor-positive, early-stage, HER2-negative invasive breast cancer who will receive adjuvant hormonal therapy for the purpose of determining whether patients can avoid adjuvant cytotoxic chemotherapy.
3. Prognosis and/or prediction of treatment response in patients with DCIS for the purpose of determining whether patients can avoid radiotherapy.
4. Prognosis and/or prediction of treatment response in patients with node-negative, early-stage, hormone receptor-positive, HER2-negative invasive breast cancer, receiving adjuvant hormonal therapy, who have survived without progression to five years post diagnosis, for the purpose of determining whether patients will continue adjuvant hormonal therapy.

For each of these indications, clinical trials have shown that there is some clinical benefit to receiving the additional therapy under consideration. However, each additional treatment has potential adverse effects. If a patient subgroup can be defined that has an extremely low risk of distant recurrence, or a subgroup can be defined that does not respond to the treatment, then the additional treatment can be forgone with little effect on cancer outcome due to the low risk of poor outcome or lack of response to treatment.

Table 3. Gene Expression Tests Reporting Recurrence Risk for Breast Cancer Considered Herein

Test	Manufacturer	Description
Oncotype DX®	Genomic Health (Redwood City, CA)	21-gene RT-PCR; identifies three groups as low, intermediate, and high risk for distant recurrence
EndoPredict®	Sividon Diagnostics (acquired by Myriad [Salt Lake City, UT] in 2016)	12-gene real-time RT-PCR; gene expression molecular score alone (EP) or EP is combined with the clinical parameters of tumor size and number positive lymph nodes (EPclin), resulting in classifications of EP low, EP high, EPclin low, or EPclin high risk for distant recurrence
Breast Cancer Index Prognostic SM	Biotheranostics (San Diego, CA)	Combines MGI and the HOXB13:IL17BR Index measured using RT-PCR; identifies two groups as low or high risk for distant recurrence
MammaPrint®	Agendia (Amsterdam, The Netherlands)	70-gene DNA microarray; identifies two groups as low or high risk for distant recurrence
Prosigna®	NanoString Technologies (Seattle, WA)	Gene expression profile is assessed by the nCounter digital platform system to determine similarity with prototypic profiles of PAM50 genes for breast cancer; identifies three categorical ROR groups (ROR-low, ROR-intermediate, ROR-high)

MGI: Molecular Grade Index; PAM50: prediction analysis of microarray 50-gene set; ROR: risk of relapse; RT-PCR: reverse transcriptase polymerase chain reaction. EP: expression profile.

Additional commercially available tests may provide some prognostic or predictive information for breast cancer. Tests intended to assess estrogen receptor, progesterone receptor, and HER2 status, such as TargetPrint® (Agendia; via quantitative microarray), are outside the scope of this review. In addition, tests that do not provide a specific recurrence risk are outside the scope of this review.

Other commercially available biomarkers are designed to provide information about tumors' molecular subtypes (i.e., luminal A, luminal B, HER2 type, and basal type). Prosigna was initially offered as a molecular subtype test. The Blueprint® 80-gene molecular subtyping assay is offered in combination with MammaPrint to augment predictive data about response to chemotherapy.

DECISION FRAMEWORK FOR EVALUATING BREAST CANCER BIOMARKERS

Simon et al Framework

Many studies have investigated individual biomarkers or combinations of biomarkers associated with breast cancer outcomes. Determining which studies constitute sufficient evidence that the test or biomarker is likely to be clinically useful depends on attributes of the test such as its performance and the quality of the study generating the results. Simon et al (2009) have described a framework to evaluate prognostic biomarker evidence.¹⁷ Study designs, such as prospective clinical trials or previously conducted clinical trials with archived tumor samples, constitute stronger evidence than studies with less planned and systematic patient recruitment and data collection. Randomized trials allow determination of treatment-biomarker interactions that may be clinically important. In some clinical scenarios, demonstration of a treatment-biomarker interaction is not critical, because the decision to withhold chemotherapy in a low-risk group (to avoid chemotherapy-related morbidity) does not require the presence of a biomarker-treatment interaction. The study must generate an absolute estimate of outcomes in the patient group of interest that would result in a change in management (e.g., withholding of chemotherapy), and the study must have sufficient precision (narrow confidence intervals). Results of the same test across studies should show the consistency of results and more than one study demonstrating the desired result should be available. Simon has proposed that at least two Simon category B studies showing results consistent with clinical utility are necessary to demonstrate adequate evidence of a biomarker.¹⁷

Breast Cancer-Specific Outcomes

The main outcome of interest for this review is 10-year distant recurrence-free survival. Distant recurrence is a hallmark of advanced breast cancer and thus more informative of overall survival than disease-free survival. Disease-free survival also includes local recurrence, which has a much better treatment prognosis than distant disease. For the extended endocrine indications in this review, the main outcome of interest is 10-year distant recurrence-free survival conditional on recurrence-free survival for five years.

Decisions to undergo or forgo adjuvant therapy (chemotherapy or endocrine) depend on how a woman values the potential benefit of lower recurrence risk relative to the harms of treatment. The balance of benefits and harms determines the thresholds that inform decisions.^{18,19} Most women will accept substantial adverse events for even modest benefit. For example, Simes et al (2001) interviewed 104 Australian women with breast cancer treated with cytotoxic chemotherapy and elicited preferences to undergo chemotherapy according to probable gain in survival.²⁰ With an expected survival of five years without chemotherapy, 73% said they would accept chemotherapy for an increased survival of six months or less; with an expected survival of 15 years, 39% would accept treatment for a gain of six months. Duric et al (2005) found 64% to 84% of 97 women expressing a willingness to undergo chemotherapy for a one year improvement in life expectancy or 3% increase in survival rates.²¹ About half felt a single day would justify adjuvant chemotherapy. A major difference between the two studies was that the chemotherapy regimen in Duric et al was less toxic. Thewes et al (2005) adopted the same approach for adjuvant endocrine therapy preferences in 102 premenopausal women with early-stage breast cancers.²² Among women having a baseline life expectancy of five years, 61% said they would accept endocrine therapy for a six month increase in life expectancy and 79% for one year; rates were similar if the baseline life expectancy was 15 years. These proportions are close to those for adjuvant chemotherapy found by Duric.

How these estimates correspond to the distant recurrence rates reported in prognostic studies is imprecise, but Henderson (2015) has suggested that below a recurrence threshold of 10% many patients will not elect adjuvant chemotherapy owing to the small absolute benefit.²³ He also noted that a majority of those patients are older with small node-negative tumors. That interpretation is consistent with a recent study of 81 women by Hamelinck et al (2016) who found that 78% of women ages 40 to 49 years, 88% ages 50 to 59, 59% ages 60 to 69, and 40% age 70 or older would accept adjuvant chemotherapy for a 0% to 10% absolute decrease in recurrence risk (see Table 4).²⁴ There was a wide range of minimally required absolute benefits, with the majority accepting chemotherapy for an absolute benefit of 1% to 5%. At a given age range, fewer women expressed a willingness to accept adjuvant endocrine therapy than chemotherapy for a given mortality benefit.

Table 4. Patient Preferences for Undergoing Adjuvant Therapy for < 10% Reduction in Recurrence Risk²⁴

Age Range, years	Proportion That Would Accept for 1% to 10% Benefit	
	Chemotherapy, %	Endocrine, %
40-49	78	78
50-59	88	44
60-69	59	63
≥ 70	40	46

Adapted from Hamelinck et al (2016).²⁴

HUMAN EPIDERMAL GROWTH FACTOR RECEPTOR 2

The human epidermal growth factor receptor (HER) family of receptor tyrosine kinases (EGFR/HER1, ErbB2/HER2, ErbB3/HER3, ErbB4/HER4) plays a major role in the pathogenesis of many solid tumors. In approximately 25% to 30% of breast cancers, overexpression of HER2 has been linked to shorter disease-free and overall survival, lack of responsiveness to tamoxifen antiestrogen therapy, and altered responsiveness to a variety of cytotoxic chemotherapy regimens.

Trastuzumab, a monoclonal antibody directed at the extracellular domain of HER2, has offered significant shorter disease-free and overall survival advantages in the metastatic and adjuvant settings in HER2-overex-

pressing patients, although not all patients respond. Fewer than 50% of patients with metastatic HER2-positive breast cancer show initial benefit from trastuzumab treatment, and many of those eventually develop resistance.¹⁻³

Current methodologies for the selection of HER2-positive patients include immunohistochemistry (IHC) to detect HER2 protein overexpression and fluorescence in situ hybridization (FISH) to detect HER2 gene amplification. However, controversy still exists regarding the accuracy, reliability, and interobserver variability of these assay methods. IHC provides a semiquantitative measure of protein levels (scored as 0, 1+, 2+, 3+) and the interpretation may be subjective. FISH is a quantitative measurement of gene amplification, in which the HER2 gene copy number is counted. However, FISH, which is considered to be more quantitative analytically, is not always representative of protein expression, and multiple studies have failed to demonstrate a relation between HER2 gene copy number and response to trastuzumab. Whereas patients who overexpress HER2 protein (IHC) or show evidence of HER2 gene amplification (FISH) have been shown to experience better outcomes on trastuzumab than those scored negative by those assays, differences in the degree of expression or amplification by these methods have generally not been shown to discriminate between groups with different outcomes. IHC and FISH testing may be affected by interlaboratory variability, and neither test provides quantitative data that reflect the activation state of signaling pathways in tumors, which may limit their utility in patient selection.⁴ Most laboratories in North America and Europe use IHC to determine HER2 protein status, with equivocal category results (2+) confirmed by FISH (or more recently by chromogenic in situ hybridization).

Typically, HER2 activates signaling pathways by dimerizing with ligand-bound epidermal growth factor receptor family members such as HER1 and HER3. A HER2 ligand has not been identified, but overexpressed HER2 is constitutively active. When HER2 is pathologically overexpressed, the receptor may homodimerize and activate signaling cascades in the absence of the normal regulatory control imposed by the requirement for ligand binding of its heterodimerization partners. A novel assay (HERmark[®] Breast Cancer Assay) was developed to quantify total HER2 protein expression and HER2 homodimers in formalin-fixed, paraffin-embedded tissue samples.

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests (LDTs) must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Oncotype DX[®], HERmark[®] Breast Cancer Assay (Monogram Biosciences) and other tests listed herein are available under the auspices of the CLIA. Laboratories that offer LDTs must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of this test.

In February 2007, MammaPrint[®] (Agendia) was cleared for marketing by the FDA through the 510(k) process for the prediction of breast cancer metastasis. In January 2015, MammaPrint[®] was cleared for marketing by the FDA through the 510(k) process for use in fresh-frozen, paraffin-embedded breast cancer tissue.

In September 2013, Prosigna[®] was cleared for marketing by the FDA through the 510(k) process. Moreover, the FDA determined that Prosigna[®] was substantially equivalent to MammaPrint[®].

FDA product code: NYI.

Currently, the Breast Cancer Index SM (Biotheranostics) and EndoPredict[®] (distributed by Myriad) are not FDA-approved.

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are

considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Colleoni M, Sun Z, Price KN, et al. Annual hazard rates of recurrence for breast cancer during 24 years of follow-up: results from the International Breast Cancer Study Group Trials I to V. *J Clin Oncol.* Mar 20 2016; 34(9):927-935. PMID 26786933
2. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 2.2017. http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed October 9, 2017.
3. Burstein HJ, Prestrud AA, Seidenfeld J, et al. American Society of Clinical Oncology clinical practice guideline: update on adjuvant endocrine therapy for women with hormone receptor-positive breast cancer. *J Clin Oncol.* Aug 10 2010;28(23):3784-3796. PMID 20625130
4. Burstein HJ, Temin S, Anderson H, et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: American Society of Clinical Oncology clinical practice guideline focused update. *J Clin Oncol.* Jul 20 2014;32(21):2255-2269. PMID 24868023
5. Wolff AC, Hammond ME, Hicks DG, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol.* Nov 1 2013;31(31):3997-4013. PMID 24101045
6. Early Breast Cancer Trialists' Collaborative Group, Davies C, Godwin J, et al. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. *Lancet.* Aug 27 2011;378(9793):771-784. PMID 21802721
7. Burstein HJ, Lacchetti C, Anderson H, et al. Adjuvant Endocrine Therapy for Women With Hormone Receptor-Positive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update on Ovarian Suppression. *J Clin Oncol.* May 10 2016;34(14):1689-1701. PMID 26884586
8. Davies C, Pan H, Godwin J, et al. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. *Lancet.* Mar 09 2013;381(9869):805-816. PMID 23219286
9. Gray RG, Rea D, Handley K, et al. aTTom: Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years in 6,953 women with early breast cancer [abstract]. *J Clin Oncol.* 2013;31(18 Suppl):5-5. PMID
10. Tormey DC, Gray R, Falkson HC. Postchemotherapy adjuvant tamoxifen therapy beyond five years in patients with lymph node-positive breast cancer. Eastern Cooperative Oncology Group. *J Natl Cancer Inst.* Dec 18 1996;88(24):1828-1833. PMID 8961972
11. Fisher B, Dignam J, Bryant J, et al. Five versus more than five years of tamoxifen for lymph node-negative breast cancer: updated findings from the National Surgical Adjuvant Breast and Bowel Project B-14 randomized trial. *J Natl Cancer Inst.* May 02 2001;93(9):684-690. PMID 11333290
12. Stewart HJ, Prescott RJ, Forrest AP. Scottish adjuvant tamoxifen trial: a randomized study updated to 15 years. *J Natl Cancer Inst.* Mar 21 2001;93(6):456-462. PMID 11259471

13. Jakesz R, Greil R, Gnant M, et al. Extended adjuvant therapy with anastrozole among postmenopausal breast cancer patients: results from the randomized Austrian Breast and Colorectal Cancer Study Group Trial 6a. *J Natl Cancer Inst.* Dec 19 2007;99(24):1845-1853. PMID 18073378
14. Goss PE, Ingle JN, Martino S, et al. A randomized trial of letrozole in postmenopausal women after five years of tamoxifen therapy for early-stage breast cancer. *N Engl J Med.* Nov 06 2003;349(19):1793-1802. PMID 14551341
15. Goss PE, Ingle JN, Martino S, et al. Randomized trial of letrozole following tamoxifen as extended adjuvant therapy in receptor-positive breast cancer: updated findings from NCIC CTG MA.17. *J Natl Cancer Inst.* Sep 07 2005;97(17):1262-1271. PMID 16145047
16. Mamounas EP, Jeong JH, Wickerham DL, et al. Benefit from exemestane as extended adjuvant therapy after 5 years of adjuvant tamoxifen: intention-to-treat analysis of the National Surgical Adjuvant Breast and Bowel Project B-33 trial. *J Clin Oncol.* Apr 20 2008;26(12):1965-1971. PMID 18332472
17. Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. *J Natl Cancer Inst.* Nov 4 2009;101(21):1446-1452. PMID 19815849
18. Pauker SG, Kassirer JP. Therapeutic decision making: a cost-benefit analysis. *N Engl J Med.* Jul 31 1975; 293(5):229-234. PMID 1143303
19. Pauker SG, Kassirer JP. The threshold approach to clinical decision making. *N Engl J Med.* May 15 1980; 302(20):1109-1117. PMID 7366635
20. Simes RJ, Coates AS. Patient preferences for adjuvant chemotherapy of early breast cancer: how much benefit is needed? *J Natl Cancer Inst Monogr.* 2001(30):146-152. PMID 11773309
21. Duric VM, Stockler MR, Heritier S, et al. Patients' preferences for adjuvant chemotherapy in early breast cancer: what makes AC and CMF worthwhile now? *Ann Oncol.* Nov 2005;16(11):1786-1794. PMID 16126738
22. Thewes B, Meiser B, Duric VM, et al. What survival benefits do premenopausal patients with early breast cancer need to make endocrine therapy worthwhile? *Lancet Oncol.* Aug 2005;6(8):581-588. PMID 16054569
23. Henderson IC. *Breast cancer: fundamentals of evidence-based disease management.* New York: Oxford University Press; 2015.
24. Hamelinck VC, Bastiaannet E, Pieterse AH, et al. A prospective comparison of younger and older patients' preferences for adjuvant chemotherapy and hormonal therapy in early breast cancer. *Clin Breast Cancer.* Oct 2016;16(5):379-388. PMID 27212474
25. Buus R, Sestak I, Kronenwett R, et al. Comparison of EndoPredict and EPclin with Oncotype DX Recurrence Score for prediction of risk of distant recurrence after endocrine therapy. *J Natl Cancer Inst.* Nov 2016; 108(11). PMID 27400969
26. Paik S, Shak S, Tang G, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med.* Dec 30 2004;351(27):2817-2826. PMID 15591335
27. Paik S, Tang G, Shak S, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol.* Aug 10 2006;24(23):3726-3734. PMID 16720680
28. Tang G, Shak S, Paik S, et al. Comparison of the prognostic and predictive utilities of the 21-gene Recurrence Score assay and Adjuvant! for women with node-negative, ER-positive breast cancer: results from NSABP B-14 and NSABP B-20. *Breast Cancer Res Treat.* May 2011;127(1):133-142. PMID 21221771
29. Sparano JA, Gray RJ, Makower DF, et al. Prospective validation of a 21-gene expression assay in breast cancer. *N Engl J Med.* Nov 19 2015;373(21):2005-2014. PMID 26412349
30. Filipits M, Rudas M, Jakesz R, et al. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin Cancer Res.* Sep 15 2011;17(18):6012-6020. PMID 21807638
31. Sgroi DC, Sestak I, Cuzick J, et al. Prediction of late distant recurrence in patients with oestrogen-receptor positive breast cancer: a prospective comparison of the breast-cancer index (BCI) assay, 21-gene recurrence score, and IHC4 in the TransATAC study population. *Lancet Oncol.* Oct 2013;14(11):1067-1076. PMID 24035531

32. Zhang Y, Schnabel CA, Schroeder BE, et al. Breast Cancer Index identifies early-stage estrogen receptor-positive breast cancer patients at risk for early- and late-distant recurrence. *Clin Cancer Res*. Aug 1 2013; 19(15):4196-4205. PMID 23757354
33. Bueno-de-Mesquita JM, Sonke GS, van de Vijver MJ, et al. Additional value and potential use of the 70-gene prognosis signature in node-negative breast cancer in daily clinical practice. *Ann Oncol*. Sep 2011;22(9): 2021-2030. PMID 19955335
34. van't Veer LJ, Yau C, Yu NY, et al. Tamoxifen therapy benefit for patients with 70-gene signature high and low risk. *Breast Cancer Res Treat*. Aug 04 2017. PMID 28776283
35. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-gene signature as an aid to treatment decisions in early-stage breast cancer. *N Engl J Med*. Aug 25 2016;375(8):717-729. PMID 27557300
36. Dowsett M, Sestak I, Lopez-Knowles E, et al. Comparison of PAM50 risk of recurrence score with Oncotype DX and IHC4 for predicting risk of distant recurrence after endocrine therapy. *J Clin Oncol*. Aug 1 2013; 31(22):2783-2790. PMID 23816962
37. Gnant M, Filipits M, Greil R, et al. Predicting distant recurrence in receptor-positive breast cancer patients with limited clinicopathological risk: using the PAM50 Risk of Recurrence score in 1478 postmenopausal patients of the ABCSG-8 trial treated with adjuvant endocrine therapy alone. *Ann Oncol*. Feb 2014;25(2): 339-345. PMID 24347518
38. Albain KS, Barlow WE, Shak S, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol*. Jan 2010;11(1):55-65. PMID 20005174
39. Dowsett M, Cuzick J, Wale C, et al. Prediction of risk of distant recurrence using the 21-gene recurrence score in node-negative and node-positive postmenopausal patients with breast cancer treated with anastrozole or tamoxifen: a TransATAC study. *J Clin Oncol*. Apr 10 2010;28(11):1829-1834. PMID 20212256
40. Gnant M, Sestak I, Filipits M, et al. Identifying clinically relevant prognostic subgroups of postmenopausal women with node-positive hormone receptor-positive early-stage breast cancer treated with endocrine therapy: a combined analysis of ABCSG-8 and ATAC using the PAM50 risk of recurrence score and intrinsic subtype. *Ann Oncol*. Aug 2015;26(8):1685-1691. PMID 25935792
41. Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat*. Jun 29 2017. PMID 28664507
42. Jasem J, Fisher CM, Amini A, et al. The 21-Gene Recurrence Score assay for node-positive, early-stage breast cancer and impact of RxPONDER Trial on chemotherapy decision-making: have clinicians already decided? *J Natl Compr Canc Netw*. Apr 2017;15(4):494-503. PMID 28404760
43. Roberts MC, Miller DP, Shak S, et al. Breast cancer-specific survival in patients with lymph node-positive hormone receptor-positive invasive breast cancer and Oncotype DX Recurrence Score results in the SEER database. *Breast Cancer Res Treat*. Jun 2017;163(2):303-310. PMID 28243896
44. Harris LN, Ismaila N, McShane LM, et al. Use of biomarkers to guide decisions on adjuvant systemic therapy for women with early-stage invasive breast cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol*. Apr 1 2016;34(10):1134-1150. PMID 26858339
45. Mook S, Schmidt MK, Viale G, et al. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1-3 positive lymph nodes in an independent validation study. *Breast Cancer Res Treat*. Jul 2009;116(2):295-302. PMID 18661261
46. Solin LJ, Gray R, Baehner FL, et al. A multigene expression assay to predict local recurrence risk for ductal carcinoma in situ of the breast. *J Natl Cancer Inst*. May 15 2013;105(10):701-710. PMID 23641039
47. Rakovitch E, Nofech-Mozes S, Hanna W, et al. A population-based validation study of the DCIS Score predicting recurrence risk in individuals treated by breast-conserving surgery alone. *Breast Cancer Res Treat*. Jul 2015;152(2):389-398. PMID 26119102
48. Esserman L, Gallant E, Alvarado M. Less is more: The evolving surgical approach to breast cancer. *Am Soc Clin Oncol Educ Book*. 2016;35:e5-e10. PMID 27249759

49. Dubsy P, Brase JC, Jakesz R, et al. The EndoPredict score provides prognostic information on late distant metastases in ER+/HER2- breast cancer patients. *Br J Cancer*. Dec 10 2013;109(12):2959-2964. PMID 24157828
50. Filipits M, Nielsen TO, Rudas M, et al. The PAM50 risk-of-recurrence score predicts risk for late distant recurrence after endocrine therapy in postmenopausal women with endocrine-responsive early breast cancer. *Clin Cancer Res*. Mar 1 2014;20(5):1298-1305. PMID 24520097
51. Sestak I, Cuzick J, Dowsett M, et al. Prediction of late distant recurrence after 5 years of endocrine treatment: a combined analysis of patients from the Austrian Breast and Colorectal Cancer Study group 8 and arimidex, tamoxifen alone or in combination randomized trials using the PAM50 risk of recurrence score. *J Clin Oncol*. Mar 10 2015;33(8):916-922. PMID 25332252
52. Sestak I, Dowsett M, Zabaglo L, et al. Factors predicting late recurrence for estrogen receptor-positive breast cancer. *J Natl Cancer Inst*. Oct 2 2013;105(19):1504-1511. PMID 24029245
53. Esserman LJ, Yau C, Thompson CK, et al. Use of molecular tools to identify patients with indolent breast cancers with ultralow risk over 2 decades. *JAMA Oncol*. Jun 29 2017. PMID 28662222
54. Sgroi DC, Carney E, Zarrella E, et al. Prediction of late disease recurrence and extended adjuvant letrozole benefit by the HOXB13/IL17BR biomarker. *J Natl Cancer Inst*. Jul 17 2013;105(14):1036-1042. PMID 23812955
55. Schroeder B, Zhang Y, Stal O, et al. Risk stratification with Breast Cancer Index for late distant recurrence in patients with clinically low-risk (T1N0) estrogen receptor-positive breast cancer. *NPJ Breast Cancer*. 2017; 3:28. PMID 28795152
56. Bosl A, Spitzmuller A, Jasarevic Z, et al. MammaPrint versus EndoPredict: Poor correlation in disease recurrence risk classification of hormone receptor positive breast cancer. *PLoS One*. 2017;12(8):e0183458. PMID 28850621
57. Sestak I, Zhang Y, Schroeder BE, et al. Cross-stratification and differential risk by Breast Cancer Index and Recurrence Score in women with hormone receptor-positive lymph node-negative early-stage breast cancer. *Clin Cancer Res*. Oct 15 2016;22(20):5043-5048. PMID 27252417
58. Hornberger J, Alvarado MD, Rebecca C, et al. Clinical validity/utility, change in practice patterns, and economic implications of risk stratifiers to predict outcomes for early-stage breast cancer: a systematic review. *J Natl Cancer Inst*. Jul 18 2012;104(14):1068-1079. PMID 22767204
59. Fan C, Oh DS, Wessels L, et al. Concordance among gene-expression-based predictors for breast cancer. *N Engl J Med*. Aug 10 2006;355(6):560-569. PMID 16899776
60. Espinosa E, Vara JA, Redondo A, et al. Breast cancer prognosis determined by gene expression profiling: a quantitative reverse transcriptase polymerase chain reaction study. *J Clin Oncol*. Oct 10 2005;23(29):7278-7285. PMID 16129846
61. Kelly CM, Bernard PS, Krishnamurthy S, et al. Agreement in risk prediction between the 21-gene recurrence score assay (Oncotype DX(R)) and the PAM50 breast cancer intrinsic Classifier in early-stage estrogen receptor-positive breast cancer. *Oncologist*. Mar 2012;17(4):492-498. PMID 22418568
62. Prat A, Parker JS, Fan C, et al. Concordance among gene expression-based predictors for ER-positive breast cancer treated with adjuvant tamoxifen. *Ann Oncol*. Nov 2012;23(11):2866-2873. PMID 22532584
63. Badve SS, Baehner FL, Gray RP, et al. Estrogen- and progesterone-receptor status in ECOG 2197: comparison of immunohistochemistry by local and central laboratories and quantitative reverse transcription polymerase chain reaction by central laboratory. *J Clin Oncol*. May 20 2008;26(15):2473-2481. PMID 18487567
64. Khoury T, Yan L, Liu S, et al. Oncotype DX RT-qPCR assay for ER and PR correlation with IHC: a study of 3 different clones. *Appl Immunohistochem Mol Morphol*. Mar 2015;23(3):178-187. PMID 24992175
65. Drukker CA, Elias SG, Nijenhuis MV, et al. Gene expression profiling to predict the risk of locoregional recurrence in breast cancer: a pooled analysis. *Breast Cancer Res Treat*. Dec 2014;148(3):599-613. PMID 25414025

66. Fitzal F, Filipits M, Rudas M, et al. The genomic expression test EndoPredict is a prognostic tool for identifying risk of local recurrence in postmenopausal endocrine receptor-positive, her2neu-negative breast cancer patients randomised within the prospective ABCSG 8 trial. *Br J Cancer*. Apr 14 2015;112(8):1405-1410. PMID 25867274
67. Krop I, Ismaila N, Stearns V. Use of biomarkers to guide decisions on adjuvant systemic therapy for women with early-stage invasive breast cancer: American Society of Clinical Oncology clinical practice focused update guideline summary. *J Oncol Pract*. Jul 11 2017;JOP2017024646. PMID 28696818
68. Duffy MJ, Harbeck N, Nap M, et al. Clinical use of biomarkers in breast cancer: Updated guidelines from the European Group on Tumor Markers (EGTM). *Eur J Cancer*. Apr 2017;75:284-298. PMID 28259011
69. Coates AS, Winer EP, Goldhirsch A, et al. Tailoring therapies—improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. *Ann Oncol*. Aug 2015;26(8):1533-1546. PMID 25939896
70. Centers for Medicare & Medicaid Services. Local Coverage Determination (LCD): MoLDX: Breast Cancer IndexSM Genetic Assay (L35631). 2015; <https://med.noridianmedicare.com/documents/10546/6990983/MoLDx+Breast+Cancer+Index+Genetic+Assay+LCD/a9495a31-0ba2-4c7b-81e4-8d424689d2c7>. Accessed October 9, 2017.
71. Zujewski JA, Kamin L. Trial assessing individualized options for treatment for breast cancer: the TAILORx trial. *Future Oncol*. Oct 2008;4(5):603-610. PMID 18922117
72. Arteaga CL, O'Neill A, Moulder SL, et al. A phase I-II study of combined blockade of the ErbB receptor network with trastuzumab and gefitinib in patients with HER2 (ErbB2)-overexpressing metastatic breast cancer. *Clin Cancer Res*. Oct 01 2008;14(19):6277-6283. PMID 18829509
73. de Alava E, Ocana A, Abad M, et al. Neuregulin expression modulates clinical response to trastuzumab in patients with metastatic breast cancer. *J Clin Oncol*. Jul 01 2007;25(19):2656-2663. PMID 17602072
74. Piccart M, Lohrisch C, Di Leo A, et al. The predictive value of HER2 in breast cancer. *Oncology*. 2001;61 Suppl 2:73-82. PMID 11694791
75. Shi Y, Huang W, Tan Y, et al. A novel proximity assay for the detection of proteins and protein complexes: quantitation of HER1 and HER2 total protein expression and homodimerization in formalin-fixed, paraffin-embedded cell lines and breast cancer tissue. *Diagn Mol Pathol*. Mar 2009;18(1):11-21. PMID 19214113
76. Monogram Biosciences. HERmark Breast Cancer Assay. 2017; <https://www.monogrambio.com/oncology-tests>. Accessed October 24, 2017.
77. Larson JS, Goodman LJ, Tan Y, et al. Analytical validation of a highly quantitative, sensitive, accurate, and reproducible assay (HERmark) for the measurement of HER2 total protein and HER2 homodimers in FFPE breast cancer tumor specimens. *Patholog Res Int*. Jun 28 2010;2010:814176. PMID 21151530
78. Huang W, Reinholz M, Weidler J, et al. Comparison of central HER2 testing with quantitative total HER2 expression and HER2 homodimer measurements using a novel proximity-based assay. *Am J Clin Pathol*. Aug 2010;134(2):303-311. PMID 20660336
79. Wolff AC, Hammond ME, Schwartz JN, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *Arch Pathol Lab Med*. Jan 2007;131(1):18-43. PMID 19548375
80. Yardley DA, Kaufman PA, Huang W, et al. Quantitative measurement of HER2 expression in breast cancers: comparison with 'real-world' routine HER2 testing in a multicenter Collaborative Biomarker Study and correlation with overall survival. *Breast Cancer Res*. Mar 18 2015;17:41. PMID 25886996
81. Bates M, Sperinde J, Kostler WJ, et al. Identification of a subpopulation of metastatic breast cancer patients with very high HER2 expression levels and possible resistance to trastuzumab. *Ann Oncol*. Sep 2011;22(9):2014-2020. PMID 21289364
82. Joensuu H, Sperinde J, Leinonen M, et al. Very high quantitative tumor HER2 content and outcome in early breast cancer. *Ann Oncol*. Sep 2011;22(9):2007-2013. PMID 21285132

83. Toi M, Sperinde J, Huang W, et al. Differential survival following trastuzumab treatment based on quantitative HER2 expression and HER2 homodimers in a clinic-based cohort of patients with metastatic breast cancer. *BMC Cancer*. Feb 23 2010;10:56. PMID 20178580
84. Lipton A, Kostler WJ, Leitzel K, et al. Quantitative HER2 protein levels predict outcome in fluorescence in situ hybridization-positive patients with metastatic breast cancer treated with trastuzumab. *Cancer*. Nov 15 2010;116(22):5168-5178. PMID 20661914
85. Lipton A, Goodman L, Leitzel K, et al. HER3, p95HER2, and HER2 protein expression levels define multiple subtypes of HER2-positive metastatic breast cancer. *Breast Cancer Res Treat*. Aug 2013;141(1):43-53. PMID 23959396
86. Han SW, Cha Y, Paquet A, et al. Correlation of HER2, p95HER2 and HER3 expression and treatment outcome of lapatinib plus capecitabine in her2-positive metastatic breast cancer. *PLoS One*. 2012;7(7):e39943. PMID 22848366
87. Duchnowska R, Biernat W, Szostakiewicz B, et al. Correlation between quantitative HER-2 protein expression and risk for brain metastases in HER-2+ advanced breast cancer patients receiving trastuzumab-containing therapy. *Oncologist*. Jan 2012;17(1):26-35. PMID 22234627
88. Barros FF, Abdel-Fatah TM, Moseley P, et al. Characterisation of HER heterodimers in breast cancer using in situ proximity ligation assay. *Breast Cancer Res Treat*. Apr 2014;144(2):273-285. PMID 24557338
89. Duchnowska R, Sperinde J, Chenna A, et al. Quantitative measurements of tumoral p95HER2 protein expression in metastatic breast cancer patients treated with trastuzumab: independent validation of the p95HER2 clinical cutoff. *Clin Cancer Res*. May 15 2014;20(10):2805-2813. PMID 24668646
90. Camp RL, Dolled-Filhart M, Rimm DL. X-tile: a new bio-informatics tool for biomarker assessment and outcome-based cut-point optimization. *Clin Cancer Res*. Nov 1 2004;10(21):7252-7259. PMID 15534099
91. Green AR, Barros FF, Abdel-Fatah TM, et al. HER2/HER3 heterodimers and p21 expression are capable of predicting adjuvant trastuzumab response in HER2+ breast cancer. *Breast Cancer Res Treat*. May 2014; 145(1):33-44. PMID 24706169
92. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: Breast cancer. Version 2.2017. http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed October 31, 2017.
93. Palmetto GBA®. MoIDX: HERmark Assay by Monogram Coding and Billing Guidelines (M00028, V10). 2017; <http://www.palmettogba.com/palmetto/MoIDX.nsf/DocsCat/MoIdx%20Website~MoIdx~Browse%20By%20Topic~Covered%20Tests~8TVSBJ3016?open&navmenu=Browse%5EBy%5ETopic%7C%7C%7C%7C>. Accessed November 15, 2017.
94. National Government Services, Inc. (Primary Geographic Jurisdiction 06 & K - Illinois, Minnesota, Wisconsin, Connecticut, New York - Entire State, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont) Local Coverage Determination (LCD): Molecular Pathology Procedures (L35000), Revision Effective Date For services performed on or after 01/01/2018.
95. Noridian Healthcare Solutions, LLC, (Jurisdiction E - California - Entire State, American Samoa, Guam, Hawaii, Northern Mariana Islands, Nevada) Local Coverage Determination (LCD): MoIDX: ONCOTYPE DX® BREAST Cancer for DCIS (Genomic Health™) (L36941), Revision Effective Date for services performed on or after 08/18/2017.
96. Palmetto GBA (Jurisdiction J & M: Alabama, Georgia, Tennessee, North Carolina, South Carolina, West Virginia, Virginia) Local Coverage Determination (LCD): MoIDX: EndoPredict BREAST Cancer Gene Expression Test (L37264), Revision Effective Date for services performed on or after 02/26/2018.
97. Palmetto GBA (Jurisdiction J & M: Alabama, Georgia, Tennessee, North Carolina, South Carolina, West Virginia, Virginia) Local Coverage Determination (LCD): MoIDX: BREAST CANCER INDEXSM Genetic Assay (L35631), Revision Effective Date for services performed on or after 02/26/2018.