

# Protocol

## Multigene Expression Assay for Predicting Recurrence in Colon Cancer

(20461)

<b>Medical Benefit</b>		<b>Effective Date:</b> 10/01/16	<b>Next Review Date:</b> 07/19
<b>Preauthorization</b>	No	<b>Review Dates:</b> 09/10, 07/11, 07/12, 07/13, 07/14, 07/15, 07/16, 07/17, 07/18	

***This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.***

*The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.*

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"><li>• With stage II or III colon cancer</li></ul>	Interventions of interest are: <ul style="list-style-type: none"><li>• Gene expression profile testing</li></ul>	Comparators of interest are: <ul style="list-style-type: none"><li>• Risk prediction based on clinicopathologic factors</li></ul>	Relevant outcomes include: <ul style="list-style-type: none"><li>• Disease-specific survival</li><li>• Test accuracy</li><li>• Test validity</li><li>• Change in disease status</li></ul>

### DESCRIPTION

Gene expression profiling (GEP) tests have been developed for use as prognostic markers in stage II or III colon cancer to help identify patients who are at high risk for recurrent disease and could be candidates for adjuvant chemotherapy.

### SUMMARY OF EVIDENCE

For individuals who have stage II or III colon cancer who receive GEP testing, the evidence includes development and validation studies and decision-impact studies. Relevant outcomes are disease-specific survival, test accuracy and validity, and change in disease status. The available evidence has shown that GEP testing for colon cancer can improve risk prediction, particularly the risk of recurrence in patients with stage II or III colon cancer. However, the degree of difference in risk conferred by the test is small. Evidence to date is insufficient to permit conclusions on whether GEP classification is sufficient to modify treatment decisions in stage II or III patients. Studies showing management changes as a consequence of testing do not demonstrate whether such changes improve outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

### POLICY

Gene expression assays for determining the prognosis of stage II or III colon cancer following surgery are considered **investigational**.

## POLICY GUIDELINES

### GENETICS NOMENCLATURE UPDATE

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing protocol updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the HUMAN Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology—"pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign"—to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely Pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

### GENETIC COUNSELING

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

### MEDICARE ADVANTAGE

The Oncotype DX® colon cancer assay may be considered **medically necessary** when used to determine prognosis and determine the treatment plan.

### BACKGROUND

Of patients with stage II colon cancer, 75% to 80% are cured by surgery alone, and the absolute benefit of

chemotherapy for the overall patient population is small. Patients most likely to benefit from chemotherapy are difficult to identify by standard clinical and pathologic risk factors. Genomic tests are intended to facilitate identifying stage II patients most likely to experience recurrence after surgery and most likely to benefit from additional treatment.

Colorectal cancer is classified as stage II (also called Dukes B) when it has spread outside the colon and/or rectum to nearby tissue but is not detectable in lymph nodes (stage III disease, also called Dukes C) and has not metastasized to distant sites (stage IV disease). Primary treatment is surgical resection of primary cancer and colonic anastomosis. After surgery, the prognosis is good, with survival rates of 75% to 80% at five years.<sup>1</sup> A 2008 meta-analysis of 50 studies of adjuvant therapy vs. surgery alone in stage II patients found a statistically significant, though small, absolute benefit of chemotherapy for disease-free survival but not for overall survival.<sup>1</sup> Therefore, adjuvant chemotherapy with 5-fluorouracil or capecitabine is recommended only for resected patients with high-risk stage II disease (i.e., those with poor prognostic features).<sup>2</sup>

However, the clinical and pathologic features used to identify high-risk disease are not well-established, and patients for whom benefits of adjuvant chemotherapy would most likely outweigh harms cannot be identified with certainty. The current diagnostic system relies on a variety of factors, including tumor substage IIB (T4A tumors that invade the muscularis propria and extend into pericorectal tissues) or IIC (T4B tumors that invade or are adherent to other organs or structures), obstruction or bowel perforation at initial diagnosis, an inadequately low number of sampled lymph nodes at surgery ( $\leq 12$ ), histologic features of aggressiveness, a high preoperative carcinoembryonic antigen level, and indeterminate or positive resection margins.<sup>2</sup>

Of interest, a 2010 review has noted that microsatellite instability and mismatch repair deficiency in colon cancer may represent confounding factors to be considered in treatment.<sup>3</sup> These factors may identify a minority (15%-20%) of the population with improved disease-free survival who may derive no benefit or may exhibit deleterious effects from adjuvant 5-fluorouracil plus leucovorin-based treatments. Patient microsatellite instability and mismatch repair status may be critically important in how to study, interpret, and use a particular gene expression profiling test.

## REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests (LDTs) must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Multigene expression assay testing for predicting recurrent colon cancer is available under the auspices of CLIA. Laboratories that offer LDTs must be licensed by the CLIA or high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Gene expression profiling tests for colon cancer currently commercially available include:

- ColoPrint® 18-Gene Colon Cancer Recurrence Assay (Agendia)
- GeneFx™ Colon (Helomics Therapeutics; also known as ColDx, Almac Diagnostics)
- OncoDefender-CRC™ (Everist Genomics)
- Oncotype DX® Colon Recurrence Score (Genomic Health).

---

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are

considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

## REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Figueredo A, Coombes ME, Mukherjee S. Adjuvant therapy for completely resected stage II colon cancer. *Cochrane Database Syst Rev.* Jul 16 2008(3):CD005390. PMID 18646127
2. National Comprehensive Cancer Network (NCCN). Clinical practice guidelines in oncology: colon cancer. Version 2.2017. [http://www.nccn.org/professionals/physician\\_gls/pdf/colon.pdf](http://www.nccn.org/professionals/physician_gls/pdf/colon.pdf). Accessed July 10, 2017.
3. Vilar E, Gruber SB. Microsatellite instability in colorectal cancer-the stable evidence. *Nat Rev Clin Oncol.* Mar 2010;7(3):153-162. PMID 20142816
4. Wang Y, Jatko T, Zhang Y, et al. Gene expression profiles and molecular markers to predict recurrence of Dukes' B colon cancer. *J Clin Oncol.* May 1 2004;22(9):1564-1571. PMID 15051756
5. Eschrich S, Yang I, Bloom G, et al. Molecular staging for survival prediction of colorectal cancer patients. *J Clin Oncol.* May 20 2005;23(15):3526-3535. PMID 15908663
6. Barrier A, Boelle PY, Roser F, et al. Stage II colon cancer prognosis prediction by tumor gene expression profiling. *J Clin Oncol.* Oct 10 2006;24(29):4685-4691. PMID 16966692
7. Barrier A, Roser F, Boelle PY, et al. Prognosis of stage II colon cancer by non-neoplastic mucosa gene expression profiling. *Oncogene.* Apr 19 2007;26(18):2642-2648. PMID 17043639
8. Blum C, Graham A, Yousefzadeh M, et al. The expression ratio of Map7/B2M is prognostic for survival in patients with stage II colon cancer. *Int J Oncol.* Sep 2008;33(3):579-584. PMID 18695889
9. O'Connell MJ, Lavery I, Yothers G, et al. Relationship between tumor gene expression and recurrence in four independent studies of patients with stage II/III colon cancer treated with surgery alone or surgery plus adjuvant fluorouracil plus leucovorin. *J Clin Oncol.* Sep 1 2010;28(25):3937-3944. PMID 20679606
10. Van Laar RK. An online gene expression assay for determining adjuvant therapy eligibility in patients with stage 2 or 3 colon cancer. *Br J Cancer.* Dec 7 2010;103(12):1852-1857. PMID 21119668
11. Wan YW, Qian Y, Rathnagiriswaran S, et al. A breast cancer prognostic signature predicts clinical outcomes in multiple tumor types. *Oncol Rep.* Aug 2010;24(2):489-494. PMID 20596637
12. Smith JJ, Deane NG, Wu F, et al. Experimentally derived metastasis gene expression profile predicts recurrence and death in patients with colon cancer. *Gastroenterology.* Mar 2010;138(3):958-968. PMID 19914252
13. Mettu RK, Wan YW, Habermann JK, et al. A 12-gene genomic instability signature predicts clinical outcomes in multiple cancer types. *Int J Biol Markers.* Oct-Dec 2010;25(4):219-228. PMID 21161944
14. Hong Y, Downey T, Eu KW, et al. A 'metastasis-prone' signature for early-stage mismatch-repair proficient sporadic colorectal cancer patients and its implications for possible therapeutics. *Clin Exp Metastasis.* Feb 2010;27(2):83-90. PMID 20143136
15. Kennedy RD, Bylesjo M, Kerr P, et al. Development and independent validation of a prognostic assay for stage II colon cancer using formalin-fixed paraffin-embedded tissue. *J Clin Oncol.* Dec 10 2011;29(35):4620-4626. PMID 22067406
16. Salazar R, Roepman P, Capella G, et al. Gene expression signature to improve prognosis prediction of stage II and III colorectal cancer. *J Clin Oncol.* Jan 1 2011;29(1):17-24. PMID 21098318

17. Marisa L, de Reynies A, Duval A, et al. Gene expression classification of colon cancer into molecular subtypes: characterization, validation, and prognostic value. *PLoS Med.* 2013;10(5):e1001453. PMID 23700391
18. Herrera M, Islam AB, Herrera A, et al. Functional heterogeneity of cancer-associated fibroblasts from human colon tumors shows specific prognostic gene expression signature. *Clin Cancer Res.* Nov 1 2013;19(21):5914-5926. PMID 24052018
19. Zhang JX, Song W, Chen ZH, et al. Prognostic and predictive value of a microRNA signature in stage II colon cancer: a microRNA expression analysis. *Lancet Oncol.* Dec 2013;14(13):1295-1306. PMID 24239208
20. Salazar R, Rosenberg R, Lutke Holzik M, et al. The PARSC trial, a prospective study for the assessment of recurrence risk in stage II colon cancer (CC) patients using ColoPrint [abstract]. *J Clin Oncol.* 2011; 29(Suppl):TPS167. PMID
21. Maak M, Simon I, Nitsche U, et al. Independent validation of a prognostic genomic signature (ColoPrint) for patients with stage II colon cancer. *Ann Surg.* Jun 2013;257(6):1053-1058. PMID 23295318
22. Kopetz S, Tabernero J, Rosenberg R, et al. Genomic classifier ColoPrint predicts recurrence in stage II colorectal cancer patients more accurately than clinical factors. *Oncologist.* Feb 2015;20(2):127-133. PMID 25561511
23. Niedzwiecki D, Frankel WL, Venook AP, et al. Association between results of a gene expression signature assay and recurrence-free interval in patients with stage II colon cancer in Cancer and Leukemia Group B 9581 (Alliance). *J Clin Oncol.* Sep 01 2016;34(25):3047-3053. PMID 27432924
24. Lenehan PF, Boardman LA, Riegert-Johnson D, et al. Generation and external validation of a tumor-derived 5-gene prognostic signature for recurrence of lymph node-negative, invasive colorectal carcinoma. *Cancer.* May 17 2012;118(21):5234-5244. PMID 22605513
25. Gray RG, Quirke P, Handley K, et al. Validation study of a quantitative multigene reverse transcriptase-polymerase chain reaction assay for assessment of recurrence risk in patients with stage II colon cancer. *J Clin Oncol.* Dec 10 2011;29(35):4611-4619. PMID 22067390
26. Venook AP, Niedzwiecki D, and Lopatin M, et al. Biologic determinants of tumor recurrence in stage II colon cancer: validation study of the 12-gene recurrence score in cancer and leukemia group B (CALGB) 9581. *J Clin Oncol.* May 10 2013;31(14):1775-1781. PMID 23530100
27. Yothers G, O'Connell MJ, Lee M, et al. Validation of the 12-gene colon cancer recurrence score in NSABP C-07 as a predictor of recurrence in patients with stage II and III colon cancer treated with fluorouracil and leucovorin (FU/LV) and FU/LV plus oxaliplatin. *J Clin Oncol.* Dec 20 2013;31(36):4512-4519. PMID 24220557
28. Reimers MS, Kuppen PJ, Lee M, et al. Validation of the 12-gene colon cancer recurrence score as a predictor of recurrence risk in stage II and III rectal cancer patients. *J Natl Cancer Inst.* Nov 2014;106(11). PMID 25261968
29. Yamanaka T, Oki E, Yamazaki K, et al. 12-gene recurrence score assay stratifies the recurrence risk in stage II/III colon cancer with surgery alone: The SUNRISE Study. *J Clin Oncol.* Aug 20 2016;34(24):2906-2913. PMID 27325854
30. Kapiteijn E, Marijnen CA, Nagtegaal ID, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med.* Aug 30 2001;345(9):638-646. PMID 11547717
31. Black ER, Falzon L, Aronson N. Gene Expression Profiling for Predicting Outcomes in Stage II Colon Cancer (Technical Brief. No. 13). Rockville, MD: Agency for Healthcare Research and Quality; 2012.
32. Brenner B, Geva R, Rothney M, et al. Impact of the 12-gene colon cancer assay on clinical decision making for adjuvant therapy in stage II colon cancer patients. *Value Health.* Jan 2016;19(1):82-87. PMID 26797240
33. Cartwright T, Chao C, Lee M, et al. Effect of the 12-gene colon cancer assay results on adjuvant treatment recommendations in patients with stage II colon cancer. *Curr Med Res Opin.* Feb 2014;30(2):321-328. PMID 24127781
34. Srivastava G, Renfro LA, Behrens RJ, et al. Prospective multicenter study of the impact of oncotype DX colon cancer assay results on treatment recommendations in stage II colon cancer patients. *Oncologist.* May 2014; 19(5):492-497. PMID 24710310

35. Noridian Healthcare Solutions, LLC, (Jurisdiction - California - Entire State, American Samoa, Guam, Hawaii, Northern Mariana Islands, Nevada) Local Coverage Determination (LCD): MoIDX: Molecular Diagnostic Tests (MDT) (L35160) Revision Effective Date 01/01/2018.