

Protocol

Occipital Nerve Stimulation

(701125)

Medical Benefit		Effective Date: 01/01/11	Next Review Date: 01/20
Preauthorization	No	Review Dates: 09/10, 07/11, 07/12, 07/13, 07/14, 07/15, 01/16, 01/17, 01/18, 01/19	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none">• With migraine headache refractory to preventive medical management	Interventions of interest are: <ul style="list-style-type: none">• Occipital nerve stimulation	Comparators of interest are: <ul style="list-style-type: none">• Medication• Self-management (e.g., relaxation, exercise)	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Functional outcomes• Quality of life• Treatment-related morbidity
Individuals: <ul style="list-style-type: none">• With non-migraine headache (e.g., hemi-crania continua, cluster headaches)	Interventions of interest are: <ul style="list-style-type: none">• Occipital nerve stimulation	Comparators of interest are: <ul style="list-style-type: none">• Medication• Self-management (e.g., relaxation, exercise)	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Functional outcomes• Quality of life• Treatment-related morbidity

DESCRIPTION

Occipital nerve stimulation (ONS) delivers a small electrical charge to the occipital nerve intended to prevent migraines and other headaches in patients who have not responded to medications. The device consists of a subcutaneously implanted pulse generator (in the chest wall or abdomen) attached to extension leads that are tunneled to join electrodes placed across one or both occipital nerves at the base of the skull. Continuous or intermittent stimulation may be used.

SUMMARY OF EVIDENCE

For individuals who have migraine headaches refractory to preventive medical management who receive ONS, the evidence includes randomized controlled trials (RCTs), systematic reviews of RCTs, and observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Systematic reviews identified five sham-controlled randomized trials. Findings from pooled analyses of these RCTs were mixed. For example, compared with placebo, response rates to ONS did not differ significantly but did reduce the number of days with prolonged moderate-to-severe headache. ONS was also associated with a substantial number of minor and serious adverse events. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have non-migraine headaches (e.g., hemicrania continua, cluster headaches) who receive ONS, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Many of the case series had small sample sizes; series with over 25 patients were available only for treatment of cluster headache. Although the case series tended to find that a substantial number of patients improved after ONS, these studies lacked blinding and comparison groups. RCTs are needed to compare outcomes between ONS and comparators (e.g., to control for a potential placebo effect). The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Occipital nerve stimulation is considered **investigational** for all indications.

BACKGROUND

HEADACHE

There are four types of headache: vascular, muscle contraction (tension), traction, and inflammatory. Primary (not the result of another condition) chronic headache is defined as headache occurring more than 15 days of the month for at least three consecutive months. An estimated 45 million Americans experience chronic headaches. For at least half of these people, the problem is severe and sometimes disabling. Herein, we only discuss types of vascular headache, including migraine, hemicrania continua, and cluster.

Migraine

Migraine is the most common type of vascular headache. Migraine headaches are usually characterized by severe pain on one or both sides of the head, an upset stomach, and, at times, disturbed vision. One year prevalence of migraine ranges from 6% to 15% in adult men and from 14% to 35% in adult women. Migraine headaches may last a day or more, and can strike as often as several times a week or as rarely as once every few years.

Treatment

Drug therapy for migraine is often combined with biofeedback and relaxation training. Sumatriptan and other triptans are commonly used for relief of symptoms. Drugs used to prevent migraine include amitriptyline, propranolol and other β -blockers, topiramate and other antiepileptic drugs, and verapamil.

Hemicrania Continua

Hemicrania continua causes moderate and occasionally severe pain on only one side of the head. At least one of the following symptoms must also occur: conjunctival injection and/or lacrimation, nasal congestion and/or rhinorrhea, or ptosis, and/or miosis. Headache occurs daily and is continuous with no pain-free periods. Hemicrania continua occurs mainly in women, and its true prevalence is not known.

Treatment

Indomethacin usually provides rapid relief of symptoms. Other nonsteroidal anti-inflammatory drugs, including ibuprofen, celecoxib, and naproxen, can provide some relief of symptoms. Amitriptyline and other tricyclic antidepressants are effective in some patients.

Cluster Headache

Cluster headache occurs in cyclical patterns or clusters of severe or very severe unilateral orbital or supraorbital and/or temporal pain. The headache is accompanied by at least one of the following autonomic symptoms: ptosis, conjunctival injection, lacrimation, rhinorrhea, and, less commonly, facial blushing, swelling, or sweating.

Bouts of one headache every other day up to eight attacks per day may last from weeks to months, usually followed by remission periods when the headache attacks stop completely. The pattern varies by person, but most people have one or two cluster periods a year. During remission, no headaches occur for months, and sometimes even years. The intense pain is caused by the dilation of blood vessels, which creates pressure on the trigeminal nerve. While this process is the immediate cause of the pain, the etiology is not fully understood. It is more common in men than in woman. One-year prevalence is estimated to be 0.5 to 1.0 in 1000.

Treatment

Management of cluster headache consists of abortive and preventive treatment. Abortive treatments include subcutaneous injection of sumatriptan, topical anesthetics sprayed into the nasal cavity, and strong coffee. Some patients respond to rapidly inhaled pure oxygen. A variety of other pharmacologic and behavioral methods of aborting and preventing attacks have been reported with wide variation in patient response.

Peripheral Nerve Stimulators

Implanted peripheral nerve stimulators have been used to treat refractory pain for many years, but have only recently been proposed to manage craniofacial pain. Occipital, supraorbital, and infraorbital stimulation have been reported in the literature.

REGULATORY STATUS

The U.S. Food and Drug Administration has not cleared or approved any occipital nerve stimulation device for treatment of headache. In 1999, the Synergy™ IPG device (Medtronic), an implantable pulse generator, was approved by the Food and Drug Administration through the premarket approval process for management of chronic, intractable pain of the trunk or limbs, and off-label use for headache is described in the literature. The Genesis™ Neuromodulation System (St. Jude Medical) was approved by the Food and Drug Administration for spinal cord stimulation and the Eon™ stimulator has received CE mark approval in Europe for the treatment of chronic migraines.

RELATED PROTOCOLS

Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy

Spinal Cord and Dorsal Root Ganglion Stimulation

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Chen YF, Bramley G, Unwin G, et al. Occipital nerve stimulation for chronic migraine--a systematic review and meta-analysis. *PLoS One*. Mar 2015;10(3):e0116786. PMID 25793740
2. Yang Y, Song M, Fan Y, et al. Occipital nerve stimulation for migraine: a systematic review. *Pain Pract*. Apr 2016;16(4):509-517. PMID 25865962
3. Saper JR, Dodick DW, Silberstein SD, et al. Occipital nerve stimulation for the treatment of intractable chronic migraine headache: ONSTIM feasibility study. *Cephalalgia*. Feb 2011;31(3):271-285. PMID 20861241
4. Silberstein SD, Dodick DW, Saper J, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: results from a randomized, multicenter, double-blinded, controlled study. *Cephalalgia*. Dec 2012;32(16):1165-1179. PMID 23034698
5. Dodick DW, Silberstein SD, Reed KL, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: long-term results from a randomized, multicenter, double-blinded, controlled study. *Cephalalgia*. Apr 2015;35(4):344-358. PMID 25078718
6. Burns B, Watkins L, Goadsby PJ. Treatment of hemicrania continua by occipital nerve stimulation with a bion device: long-term follow-up of a crossover study. *Lancet Neurol*. Nov 2008;7(11):1001-1012. PMID 18845482
7. Burns B, Watkins L, Goadsby PJ. Treatment of intractable chronic cluster headache by occipital nerve stimulation in 14 patients. *Neurology*. Jan 27 2009;72(4):341-345. PMID 19171831
8. Magis D, Gerardy PY, Remacle JM, et al. Sustained effectiveness of occipital nerve stimulation in drug-resistant chronic cluster headache. *Headache*. Sep 2011;51(8):1191-1201. PMID 21848953
9. Mueller OM, Gaul C, Katsarava Z, et al. Occipital nerve stimulation for the treatment of chronic cluster headache - lessons learned from 18 months experience. *Cen Eur Neurosurg*. May 2011;72(2):84-89. PMID 21448856
10. Fontaine D, Blond S, Lucas C, et al. Occipital nerve stimulation improves the quality of life in medically-intractable chronic cluster headache: Results of an observational prospective study. *Cephalalgia*. Oct 2017;37(12):1173-1179. PMID 27697849
11. Leone M, Proietti Cecchini A, Messina G, et al. Long-term occipital nerve stimulation for drug-resistant chronic cluster headache. *Cephalalgia*. Jul 2017;37(8):756-763. PMID 27250232
12. Miller S, Watkins L, Matharu M. Treatment of intractable chronic cluster headache by occipital nerve stimulation: a cohort of 51 patients. *Eur J Neurol*. Feb 2017;24(2):381-390. PMID 27995704
13. Vadivelu S, Bolognese P, Milhorat TH, et al. Occipital nerve stimulation for refractory headache in the Chiari malformation population. *Neurosurgery*. Jun 2012;70(6):1430-1436; discussion 1436-1437. PMID 22418582
14. Sweet JA, Mitchell LS, Narouze S, et al. Occipital nerve stimulation for the treatment of patients with medically refractory occipital neuralgia: Congress of Neurological Surgeons Systematic Review and Evidence-Based Guideline. *Neurosurgery*. Sep 2015;77(3):332-341. PMID 26125672
15. National Institute for Health and Care Excellence. Occipital nerve stimulation for intractable chronic migraine [IPG452]. 2013; <https://www.nice.org.uk/guidance/ipg452>. Accessed March 8, 2018.