

(701151)

<b>Medical Benefit</b>		<b>Effective Date:</b> 04/01/19	<b>Next Review Date:</b> 01/20
<b>Preauthorization</b>	No	<b>Review Dates:</b> 05/17, 01/18, 01/19	

**Preauthorization is not required.**

*The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.*

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> <li>With lower urinary tract obstruction symptoms due to benign prostatic hyperplasia</li> </ul>	Interventions of interest are: <ul style="list-style-type: none"> <li>Prostatic urethral lift</li> </ul>	Comparators of interest are: <ul style="list-style-type: none"> <li>Transurethral resection of the prostate</li> <li>Minimally invasive prostate resection or ablation</li> <li>Continued medical management</li> </ul>	Relevant outcomes include: <ul style="list-style-type: none"> <li>Symptoms</li> <li>Functional outcomes</li> <li>Health status measures</li> <li>Quality of life</li> <li>Treatment-related morbidity</li> </ul>

### DESCRIPTION

Benign prostatic hyperplasia (BPH) is a common condition in older individuals that can lead to increased urinary frequency, an urgency to urinate, a hesitancy to urinate, nocturia, and a weak stream when urinating. The prostatic urethral lift (PUL) procedure involves the insertion of one or more permanent implants into the prostate, which retracts prostatic tissue and maintains an expanded urethral lumen.

### SUMMARY OF EVIDENCE

For individuals who have lower urinary tract obstruction symptoms due to BPH and receive a PUL, the evidence includes systematic reviews, randomized controlled trials (RCTs), and noncomparative studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. One RCT, the BPH6 study, compared the PUL procedure with transurethral resection of the prostate and reported that the PUL procedure was noninferior for the study's composite end point, which required concurrent fulfillment of six independently validated measures of symptoms, safety, and sexual health. While transurethral resection of the prostate was superior to PUL in managing lower urinary tract symptoms, PUL did provide significant symptom improvement over two years. PUL was further superior to transurethral resection of the prostate in preserving sexual function. These findings were corroborated by another RCT (the LIFT study), which compared PUL with sham control. Patients underwent washout of BPH medications before enrollment. LIFT reported that patients with the PUL procedure, compared with patients who had sham surgery and no BPH medication, had greater improvements in lower urinary tract symptoms without worsened sexual function at three months. After three months, patients were given the option to have PUL surgery; 80% of the patients with sham procedures chose that option. Publications from this trial reported that functional improvements were

durable over three-, four-, and five-year follow-ups in a subset of patients treated with PUL; there was a high number of exclusions and loss to follow-up in that group. The evidence is sufficient to determine the effects of the technology on health outcomes.

## POLICY

Use of prostatic urethral lift in individuals with moderate to severe lower urinary tract obstruction due to benign prostatic hyperplasia may be considered **medically necessary** when all of the following criteria are met:

- The patient has persistent or progressive lower urinary tract symptoms despite medical therapy ( $\alpha_1$ -adren-  
ergic antagonists maximally titrated, 5 $\alpha$ -reductase inhibitors, or combination medication therapy maximally  
titrated) over a trial period of no less than a six months, or is unable to tolerate medical therapy; AND,
- Prostate gland volume is 80 mL or less; AND,
- Prostate anatomy demonstrates normal bladder neck without an obstructive or protruding median lobe;  
AND,
- Patient does not have urinary retention, urinary tract infection, or recent prostatitis (within past year); AND,
- Patient does not have prostate-specific antigen level of three ng/mL or higher, or has had appropriate test-  
ing to exclude diagnosis of prostate cancer; AND,
- Patient does not have a contact dermatitis nickel allergy.

Use of prostatic urethral lift in other situations is considered **investigational**.

## BACKGROUND

### BENIGN PROSTATIC HYPERPLASIA

BPH is a common disorder among older individuals that results from hyperplastic nodules in the periurethral or transitional zone of the prostate. BPH prevalence increases with age and is present in more than 80% of individuals ages 70 to 79.<sup>1</sup> The clinical manifestations of BPH include increased urinary frequency, nocturia, an urgency or hesitancy to urinate, and a weak stream when urinating. The urinary tract symptoms often progress with worsening hypertrophy and may lead to acute urinary retention, incontinence, renal insufficiency, and/or urinary tract infection.

Two scores are widely used to evaluate BPH-related symptoms: the American Urological Association Symptom Index (AUASI) and the International Prostate Symptom Score (IPSS). The AUASI is a self-administered seven item questionnaire assessing the severity of various urinary symptoms.<sup>2</sup> The IPSS incorporates questions from the AUASI and a quality of life question or a "Bother score."<sup>3</sup>

### Management

Evaluation and management of BPH include assessment for other causes of lower urinary tract dysfunction (e.g., prostate cancer), symptom severity, and the degree that symptoms are bothersome to determine the therapeutic approach.

### Medical Therapy

A discussion about medical therapy is generally indicated for patients with moderate-to-severe symptoms (e.g., an AUASI score of eight or more), bothersome symptoms, or both. Available medical therapies for BPH-related lower urinary tract dysfunction include  $\alpha$ -adrenergic blockers (e.g., alfuzosin, doxazosin, tamsulosin, terazosin, silodosin), 5 $\alpha$ -reductase inhibitors (e.g., finasteride, dutasteride), combination  $\alpha$ -adrenergic blockers and 5 $\alpha$ -

reductase inhibitors, anti-muscarinic agents (e.g., darifenacin, solifenacin, oxybutynin), and phosphodiesterase-5 inhibitors (e.g., tadalafil).<sup>1</sup> In a meta-analysis of both indirect comparisons from placebo-controlled studies (including 6333 patients) and direct comparative studies (including 507 patients), Djavan et al (1999) found that the IPSS improved by 30% to 40% and the Qmax score (mean peak urinary flow rate) improved by 16% to 25% in individuals assigned to  $\alpha$ -adrenergic blockers.<sup>4</sup> Combination therapy using an  $\alpha$ -adrenergic blocker and 5 $\alpha$ -reductase inhibitor has been shown to be more effective for improving IPSS than either treatment alone, with median scores improving by more than 40% over one year and by more than 45% over four years.<sup>5</sup>

### Surgical and Ablative Therapies

Patients who do not have sufficient response to medical therapy, or who are experiencing significant side effects with medical therapy, may be referred for surgical or ablative therapies. Various surgical and ablative procedures are used to treat BPH. Transurethral resection of the prostate is generally considered the reference standard for comparisons of BPH procedures.<sup>6</sup> In the perioperative period, transurethral resection of the prostate is associated with risks of any operative procedure (e.g., anesthesia risks, blood loss). Although short-term mortality risks are generally low, a large prospective study with 10,654 patients by Reich et al (2008) reported the following short-term complications: "failure to void (5.8%), surgical revision (5.6%), significant urinary tract infection (3.6%), bleeding requiring transfusions (2.9%), and transurethral resection syndrome (1.4%)."<sup>7</sup> Incidental carcinoma of the prostate was diagnosed by histologic examination in 9.8% of patients. In the longer term, transurethral resection of the prostate is associated with increased risk of sexual dysfunction and incontinence.

Several minimally invasive prostate ablation procedures are available, including transurethral microwave thermotherapy, transurethral needle ablation of the prostate, urethromicroablation phototherapy, and photoselective vaporization of the prostate. The minimally invasive procedures were individually compared with transurethral resection of the prostate at the time they were developed, which provided a general benchmark for evaluating those procedures.

### REGULATORY STATUS

One implantable transprostatic tissue retractor system has been cleared for marketing by the Food and Drug Administration (FDA) through the 510(k) process. In 2013, the NeoTract UroLift® System UL400 (NeoTract) was cleared (after receiving clearance through FDA's de novo classification process in March 2013; K130651/DEN130023). In 2016, FDA determined that the UL500 was substantially equivalent to existing devices (UL400) for the treatment of symptoms of urinary flow obstruction secondary to BPH in individuals ages 50 years and older. In 2017, FDA expanded the indication for the UL400 and UL500 to include lateral and median lobe hyperplasia in men 45 years or older. FDA product code: PEW.

---

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

**REFERENCES**

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Sarma AV, Wei JT. Clinical practice. Benign prostatic hyperplasia and lower urinary tract symptoms. *N Engl J Med*. Jul 19 2012;367(3):248-257. PMID 22808960
2. Barry MJ, Fowler FJ, Jr., O'Leary MP, et al. The American Urological Association symptom index for benign prostatic hyperplasia. The Measurement Committee of the American Urological Association. *J Urol*. Nov 1992;148(5):1549-1557; discussion 1564. PMID 1279218
3. O'Leary M P. Validity of the "bother score" in the evaluation and treatment of symptomatic benign prostatic hyperplasia. *Rev Urol*. Winter 2005;7(1):1-10. PMID 16985801
4. Djavan B, Marberger M. A meta-analysis on the efficacy and tolerability of alpha1-adrenoceptor antagonists in patients with lower urinary tract symptoms suggestive of benign prostatic obstruction. *Eur Urol*. Jun 1999;36(1):1-13. PMID 10364649
5. McConnell JD, Roehrborn CG, Bautista OM, et al. The long-term effect of doxazosin, finasteride, and combination therapy on the clinical progression of benign prostatic hyperplasia. *N Engl J Med*. Dec 18 2003;349(25):2387-2398. PMID 14681504
6. Foster HE, Barry MJ, Dahm P, et al. Surgical management of lower urinary tract symptoms attributed to benign prostatic hyperplasia: AUA Guideline. *J Urol*. Jun 11 2018. PMID 29775639
7. Reich O, Gratzke C, Bachmann A, et al. Morbidity, mortality and early outcome of transurethral resection of the prostate: a prospective multicenter evaluation of 10,654 patients. *J Urol*. Jul 2008;180(1):246-249. PMID 18499179
8. Sundaram D, Sankaran PK, Raghunath G, et al. Correlation of Prostate Gland Size and Uroflowmetry in Patients with Lower Urinary Tract Symptoms. *J Clin Diagn Res*. May 2017;11(5):AC01-AC04. PMID 28658743
9. Rosen RC, Catania JA, Althof SE, et al. Development and validation of four-item version of Male Sexual Health Questionnaire to assess ejaculatory dysfunction. *Urology*. May 2007;69(5):805-809. PMID 17482908
10. Cappelleri JC, Rosen RC. The Sexual Health Inventory for Men (SHIM): a 5-year review of research and clinical experience. *Int J Impot Res*. Jul-Aug 2005;17(4):307-319. PMID 15875061
11. Sonksen J, Barber NJ, Speakman MJ, et al. Prospective, randomized, multinational study of prostatic urethral lift versus transurethral resection of the prostate: 12-month results from the BPH6 study. *Eur Urol*. Oct 2015;68(4):643-652. PMID 25937539
12. Barry MJ, Williford WO, Chang Y, et al. Benign prostatic hyperplasia specific health status measures in clinical research: how much change in the American Urological Association symptom index and the benign prostatic hyperplasia impact index is perceptible to patients? *J Urol*. Nov 1995;154(5):1770-1774. PMID 7563343
13. Roehrborn CG, Wilson TH, Black LK. Quantifying the contribution of symptom improvement to satisfaction of men with moderate to severe benign prostatic hyperplasia: 4-year data from the CombAT trial. *J Urol*. May 2012;187(5):1732-1738. PMID 22425127
14. McVary KT, Roehrborn CG, Avins AL, et al. American Urological Association Guideline: Management of Benign Prostatic Hyperplasia (BPH). 2010 (affirmed 2014); [http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-\(2010-reviewed-and-validity-confirmed-2014\)](http://www.auanet.org/guidelines/benign-prostatic-hyperplasia-(2010-reviewed-and-validity-confirmed-2014)). Accessed November 7, 2017.
15. Barry MJ, Fowler FJ, Jr., O'Leary MP, et al. Measuring disease-specific health status in men with benign prostatic hyperplasia. Measurement Committee of The American Urological Association. *Med Care*. Apr 1995;33(4 Suppl):AS145-155. PMID 7536866
16. Perera M, Roberts MJ, Doi SA, et al. Prostatic urethral lift improves urinary symptoms and flow while preserving sexual function for men with benign prostatic hyperplasia: a systematic review and meta-analysis. *Eur Urol*. Apr 2015;67(4):704-713. PMID 25466940
17. Garrido Abad P, Coloma Del Peso A, Sinues Ojas B, et al. Urolift(R), a new minimally invasive treatment for patients with low urinary tract symptoms secondary to BPH. Preliminary results. *Arch Esp Urol*. Jul-Aug 2013;66(6):584-591. PMID 23985459

18. Hoffman RM, Monga M, Elliott SP, et al. Microwave thermotherapy for benign prostatic hyperplasia. *Cochrane Database Syst Rev*. Sep 12 2012;9(9):CD004135. PMID 22972068
19. Shore N, Freedman S, Gange S, et al. Prospective multi-center study elucidating patient experience after prostatic urethral lift. *Can J Urol*. Feb 2014;21(1):7094-7101. PMID 24529008
20. McNicholas TA, Woo HH, Chin PT, et al. Minimally invasive prostatic urethral lift: surgical technique and multinational experience. *Eur Urol*. Aug 2013;64(2):292-299. PMID 23357348
21. Chin PT, Bolton DM, Jack G, et al. Prostatic urethral lift: two-year results after treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Urology*. Jan 2012;79(1):5-11. PMID 22202539
22. Woo HH, Bolton DM, Laborde E, et al. Preservation of sexual function with the prostatic urethral lift: a novel treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *J Sex Med*. Feb 2012;9(2):568-575. PMID 22172161
23. Woo HH, Chin PT, McNicholas TA, et al. Safety and feasibility of the prostatic urethral lift: a novel, minimally invasive treatment for lower urinary tract symptoms (LUTS) secondary to benign prostatic hyperplasia (BPH). *BJU Int*. Jul 2011;108(1):82-88. PMID 21554526
24. Cantwell AL, Bogache WK, Richardson SF, et al. Multicentre prospective crossover study of the 'prostatic urethral lift' for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia. *BJU Int*. Apr 2014;113(4):615-622. PMID 24765680
25. Roehrborn CG, Gange SN, Shore ND, et al. The prostatic urethral lift for the treatment of lower urinary tract symptoms associated with prostate enlargement due to benign prostatic hyperplasia: the L.I.F.T. Study. *J Urol*. Dec 2013;190(6):2161-2167. PMID 23764081
26. McVary KT, Gange SN, Shore ND, et al. Treatment of LUTS secondary to BPH while preserving sexual function: randomized controlled study of prostatic urethral lift. *J Sex Med*. Jan 2014;11(1):279-287. PMID 24119101
27. Shore N. A review of the prostatic urethral lift for lower urinary tract symptoms: symptom relief, flow improvement, and preservation of sexual function in men with benign prostatic hyperplasia. *Curr Bladder Dysfunct Rep*. Mar 27 2015;10(2):186-192. PMID 25984251
28. Roehrborn CG, Rukstalis DB, Barkin J, et al. Three year results of the prostatic urethral L.I.F.T. study. *Can J Urol*. Jun 2015;22(3):7772-7782. PMID 26068624
29. Jones P, Rajkumar GN, Rai BP, et al. Medium-term outcomes of Urolift (minimum 12 months follow-up): evidence from a systematic review. *Urology*. Nov 2016;97:20-24. PMID 27208817
30. Bozkurt A, Karabakan M, Keskin E, et al. Prostatic urethral lift: a new minimally invasive treatment for lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Urol Int*. Nov 2016;96(2):202-206. PMID 26613256
31. Ray A, Morgan H, Wilkes A, et al. The Urolift System for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia: a NICE Medical Technology Guidance. *Appl Health Econ Health Policy*. Oct 2016;14(5):515-526. PMID 26832146
32. Gratzke C, Barber N, Speakman MJ, et al. Prostatic urethral lift vs. transurethral resection of the prostate: 2-year results of the BPH6 prospective, multicentre, randomized study. *BJU Int*. May 2017;119(5):767-775. PMID 27862831
33. Roehrborn CG, Barkin J, Gange SN, et al. Five year results of the prospective randomized controlled prostatic urethral L.I.F.T. study. *Can J Urol*. Jun 2017;24(3):8802-8813. PMID 28646935
34. Rukstalis D, Rashid P, Bogache WK, et al. 24-month durability after crossover to the prostatic urethral lift from randomised, blinded sham. *BJU Int*. Oct 2016;118(Suppl 3):14-22. PMID 27684483
35. Roehrborn CG. Prostatic urethral lift: a unique minimally invasive surgical treatment of male lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Urol Clin North Am*. Aug 2016;43(3):357-369. PMID 27476128
36. National Institute for Health and Care Excellence (NICE). Insertion of prostatic urethral lift implants to treat lower urinary tract symptoms secondary to benign prostatic hyperplasia [IPG475]. 2014; <https://www.nice.org.uk/guidance/ipg475/chapter/1-recommendations>. Accessed July 20, 2018.

37. National Institute for Health and Care Excellence (NICE). UroLift for treating lower urinary tract symptoms of benign prostatic hyperplasia [MTG26]. 2015; <https://www.nice.org.uk/guidance/mtg26>. Accessed July 20, 2018.